

INTERNATIONAL CENTRE FOR ECONOMIC RESEARCH



WORKING PAPER SERIES

Massimo Marinacci and Luigi Montrucchio

**SUBCALCULUS FOR SET FUNCTIONS
AND CORES OF TU GAMES**

Working Paper no. 09/2001
April 2001

**APPLIED MATHEMATICS
WORKING PAPER SERIES**



Subcalculus for Set Functions and Cores of TU Games

Massimo Marinacci and Luigi Montrucchio

Dipartimento di Statistica e Matematica Applicata

Università di Torino,

10122 Torino, Italy

and ICER

massimo@econ.unito.i

luigi.montrucchio@econ.unito.it

April 10, 2001

Abstract

This paper introduces a subcalculus for general set functions and uses this framework to study the core of TU games. After stating a linearity theorem, we establish several theorems that characterize measure games having finite-dimensional cores. This is a very tractable class of games relevant in many economic applications. Finally, we show that exact games with finite dimensional cores are generalized linear production games.

Keywords. TU games, non-additive set functions, subcalculus, cores.

1 Introduction

General set functions, not necessarily additive, are widely used in mathematical economics. In cooperative game theory, the key notion of transferable utility (TU) game is modelled as a general set function ν defined on a collection Σ of admissible coalitions, with the only requirement on ν that it

takes on value zero at the empty set. In decision theory, non-additive set functions have been recently used to model “vague” beliefs, which in general are not representable by standard additive probabilities (see Schmeidler, 1989). Though the motivation is very different, the mathematical object is essentially the same in both cases.

This has motivated a large literature on non-additive set functions in both game and decision theory, that includes the classic book of Aumann and Shapley (1974). In mathematics as well, non-additive set functions have been the subject of many investigations, mostly in the wake of the seminal work of Choquet (1953), which anticipated most of the themes of the subsequent literature.

Rather surprisingly, in these different strands of literature there has been little attempt to develop a systematic calculus and subcalculus for general set functions, despite of the potential insights that such basic mathematical tools could provide. Recently, Epstein and Marinacci (2000) have developed a calculus for TU games, in which the derivative is an additive set function that suitably approximates the TU game on “small” sets. This derivative is then used to study the core of TU games. In their analysis a key role is played by linear sets (coalitions), namely sets E in Σ such that $\nu(E) + \nu(E^c) = \nu(\Omega)$, where Ω is the grand coalition. Naturally, the empty set \emptyset and the grand coalition Ω are linear sets. They show that, under mild assumptions, the core shrinks to a singleton as long as the game is differentiable at some linear set. Moreover, the core consists of the derivative itself.

A limitation of their analysis is that the core may not be a singleton. This naturally leads to the question of whether it is possible to extend their approach by using superdifferentials rather than differentials. This is our purpose in the present work, where a subcalculus for TU games is introduced and exploited to characterize cores of TU games.

Our starting point was the discovery of a simple characterization of the cores by means of superdifferentials. As a matter of fact, let $\partial\nu(E)$ be the natural adaptation for TU games of the standard superdifferential of functions on Euclidean spaces. For the core of a TU game ν it holds

$$\text{core}(\nu) = \partial\nu(E) \cap \partial\nu(E^c),$$

where E is any linear set (Theorem 10). Based on this simple characterization we are able to prove several novel results and to provide simple proofs and a unifying framework for some important known results. In particular, our

“subcalculus” framework is the natural setting in which some of the powerful methods of Convex Analysis can be used to study TU games.

More specifically, our paper is organized as follows. In Section 3 we discuss the main properties of the superdifferentials. They turn out to be similar to those of the standard superdifferentials, though the notions are less close than one might think at a first sight. Among them, it is especially important the sum rule for convex games, which ensures that $\partial(\nu_1 + \nu_2)(E) = \partial\nu_1(E) + \partial\nu_2(E)$ for all sets E in Σ . An immediate consequence of this rule is that the cores of convex games are stable under summation, that is, $core(\nu_1 + \nu_2) = core(\nu_1) + core(\nu_2)$.

After having established a “subcalculus,” Section 4 studies the relations existing among our superdifferentials, the derivatives studied by Epstein and Marinacci (2000), and the cores. The main result, Theorem 12, provides conditions ensuring that the core shrinks to a singleton as long as the differential of the game belongs to its superdifferential. This result can be viewed as an enrichment of the theory developed by Epstein and Marinacci (2000).

In Section 5 we specialize our analysis to measure games. As a matter of fact, TU games that are relevant for economic applications have often the form $\nu = g(P)$, where $P = (P_1, \dots, P_N) : \Sigma \rightarrow \mathbb{R}^N$ is a nonatomic vector measure and $g : \mathbb{R}^N \rightarrow \mathbb{R}$ is a function such that $\nu(E) = g(P(E))$ for all sets E belonging to Σ . Games of this form are called *measure games*, and standard examples include exchange economies with transferable utilities and models of production technology. While we do not expatiate here on these known issues, we refer the reader to Aumann and Shapley (1974) and Hart and Neyman (1988) for detailed discussions of these examples and of the relevance of measure games in economic applications.

Section 6 is entirely devoted to linear measure games. Since, in general, the cores of TU games are large and difficult to describe, it is important to consider games having tractable cores. Based on our subcalculus, we provide simple conditions under which the cores of measure games consist of linear combinations $\sum_{i=1}^N \alpha_i P_i$ of the components $\{P_i\}_{i=1}^N$ of the underlying vector measure P . For example, we show that the cores have this form whenever there exists a linear and radial set $E \in \Sigma$, that is, a linear set E such that $P(E)$ belongs to the relative interior of $R(P)$, the range $\{P(E) : E \in \Sigma\} \subseteq \mathbb{R}^N$ of the vector measure P . The existence of linear and radial sets is a conditions often satisfied by economic games of the form $g(P)$. In fact, these games typically feature some homogeneity condition of the function g , and it will be seen that even very mild homogeneity conditions deliver linear and

radial sets.

Our results of this section generalize well-known results of Billera and Raanan (1981), as well as recent results of Einy, Moreno, and Shitovitz (1999). They are based on a novel linearity theorem for nonatomic vector measures (Theorem 20) that should be of independent interest. This theorem relies on results from both measure theory and convex analysis, an interplay made possible by the Lyapunov Theorem, which guarantees that the range $R(P)$ is a convex set.

In Section 7 we consider general games, not necessarily measure ones, that have finite-dimensional cores. The main result of the section shows that an exact game has a finite-dimensional core if and only if it is a generalized linear production games, a very tractable class of measure games introduced in Section 6. This result is interesting because many economic games have finite-dimensional cores (see Hart and Neyman, 1988), and our characterization shows that, when exact, all these games are nothing but generalized linear production games.

Finally, in the Concluding Remarks we discuss the related works of Billera and Raanan (1981) and Einy, Moreno, and Shitovitz (1999), as well as the relationships between linear cores and semi-infinite linear programming. The Appendix gathers some technical lemmas and lengthy proofs.

2 Preliminaries

Throughout the paper, Ω is the set of players and Σ is the σ -algebra of admissible coalitions. Subsets of Ω are understood to be in Σ even where not stated explicitly.

A set function $\nu : \Sigma \rightarrow \mathbb{R}$ is a *game* if $\nu(\emptyset) = 0$. A game ν is

positive if $\nu(E) \geq 0$ for all E ,

bounded if $\sup_{E \in \Sigma} |\nu(E)| < \infty$.

monotone if $\nu(E) \geq \nu(E')$ whenever $E' \subseteq E$,

superadditive if $\nu(E \cup E') \geq \nu(E) + \nu(E')$ for all pairwise disjoint sets E and E' ,

supermodular (or convex) if $\nu(E \cup E') + \nu(E \cap E') \geq \nu(E) + \nu(E')$ for all sets E and E' ,

additive (or a charge) if $\nu(E \cup E') = \nu(E) + \nu(E')$ for all pairwise disjoint sets E and E' ,

countably additive (or a measure) if $\nu(\bigcup_{i=1}^{\infty} E_i) = \sum_{i=1}^{\infty} \nu(E_i)$ for all countable collections of pairwise disjoint sets $\{E_i\}_{i=1}^{\infty}$.

Unless otherwise stated, charges and measures are understood to be signed. The set of all charges (measures) that are bounded with respect to the variation norm is denoted by $ba(\Omega)$ ($ca(\Omega)$). Generic elements of $ba(\Omega)$ are denoted by m , while its nonnegative elements are denoted by P .

A charge m is *non-atomic* if for all $m(E) \neq 0$ there exists $B \subseteq E$ such that $m(B) \neq 0$ and $m(E - B) \neq 0$. It is *strongly continuous* if, for every $\varepsilon > 0$, there exists a partition $\{E_1, \dots, E_n\}$ of Ω in Σ such that $|m|(E_i) \leq \varepsilon$ for all $i = 1, \dots, n$. A strongly continuous charge is non-atomic, while the converse holds only for measures (see Bhaskara Rao and Bhaskara Rao (1982)). Let $m = (m_1, \dots, m_N) : \Sigma \rightarrow \mathbb{R}^N$ be a vector charge. If each m_i is strongly continuous, then by the Lyapunov Theorem the range $R(m) = \{m(A) : A \in \Sigma\}$ is a convex subset of \mathbb{R}^N (see Bhaskara Rao and Bhaskara Rao (1982)).

The game $\nu : \Sigma \rightarrow \mathbb{R}$ is a *measure game* if there exists a positive vector charge $P = (P_1, \dots, P_N) : \Sigma \rightarrow \mathbb{R}_+^N$, with each $P_i : \Sigma \rightarrow \mathbb{R}_+$ bounded and strongly continuous, and a function $g : R(P) \rightarrow \mathbb{R}$ such that

$$\nu(E) = g(P(E)) \quad \text{for all } E \in \Sigma.$$

When $N = 1$, $\nu = g(P)$ is called a *scalar measure game*.

The *core* of a game ν is

$$\text{core}(\nu) = \{m \in ba(\Omega) : m(\Omega) = \nu(\Omega) \text{ and } m(E) \geq \nu(E) \text{ for all } E \in \Sigma\}.$$

If the game is bounded, the core is w^* -compact by the Nikodym boundedness theorem (see, e.g., Diestel (1984) p. 80). A game ν is *exact* if $\text{core}(\nu) \neq \emptyset$ and $\nu(E) = \min_{m \in \text{core}(\nu)} m(E)$ for all $E \in \Sigma$. All positive convex games are exact (see Schmeidler, 1972).

Given a game $\nu : \Sigma \rightarrow \mathbb{R}$, a set E is *linear* if $\nu(E) + \nu(E^c) = \nu(\Omega)$. Notice that both Ω and \emptyset are linear sets. Moreover, when $\text{core}(\nu) \neq \emptyset$, E is linear if and only if $\nu(E) + \nu(E^c) \geq \nu(\Omega)$. The set of linear sets is denoted by \mathcal{A} .

Linear sets are delivered by *efficient coalition structures*, that is, at most countable partitions $\{E_i\}_{i \in I}$ of Ω such that $\sum_{i \in I} \nu(E_i) = \nu(\Omega)$. In fact, if

ν is superadditive and either ν is continuous or the partition is finite, then E_i is linear for each i in I (see Epstein and Marinacci (2000)).¹

We close by reporting the notion of derivative for games studied by Epstein and Marinacci (2000). For any $A \in \Sigma$, let $\{A^{j,\lambda}\}_{j=1}^{n_\lambda}$ be a finite partition of A . Denote by $\{A^{j,\lambda}\}_\lambda$ the net of all finite partitions of A , where $\lambda' > \lambda$ implies that the partition corresponding to λ' refines that corresponding to λ .

Definition 1 *A game $\nu : \Sigma \rightarrow \mathbb{R}$ is differentiable at $E \in \Sigma$ if there exists a charge $\delta\nu(\cdot; E) \in ba(\Omega)$ such that for all $F \subseteq E^c$ and $G \subseteq E$,*

$$\sum_{j=1}^{n_\lambda} |\nu(E \cup F^{j,\lambda} - G^{j,\lambda}) - \nu(E) - \delta\nu(F^{j,\lambda}; E) + \delta\nu(G^{j,\lambda}; E)| \xrightarrow{\lambda} 0.$$

This definition is slightly different from that of Epstein and Marinacci (2000), which originates in Epstein (1999), as we do not require the charge $\delta\nu(\cdot; E)$ to be convex-ranged.

3 Superdifferentials

Definition 2 *A game $\nu : \Sigma \rightarrow \mathbb{R}$ is superdifferentiable at $E \in \Sigma$ if there exists a charge $m \in ba(\Omega)$ such that*

$$\nu(A) \leq \nu(E) + m(A) - m(E) \tag{1}$$

for each $A \in \Sigma$.

The charges m that satisfy Eq. (1) are called *supergradients* and $\partial\nu(E)$ is the *superdifferential* of ν , that is, the (possibly empty) set of all supergradients.

Definition 2 is the natural adaptation to our setting of the standard notion of superdifferential of real-valued functions (see Rockafellar, 1970),² as it becomes evident by considering measure games $g(P) : \Sigma \rightarrow \mathbb{R}$. Recall that,

¹In a finite setting, efficient coalition structures have been introduced by Aumann and Dreze (1974).

²Fujishige (1991) gives a similar definition for supermodular functions defined on finite distributive lattices.

given a subset $A \subseteq \mathbb{R}^N$ (e.g., $A = R(P)$), a function $g : A \rightarrow \mathbb{R}$ is *superdifferentiable* at $x_0 \in A$ if there is a vector $\chi \in \mathbb{R}^N$, called *supergradient*, such that $g(x_0) \leq g(x) + \chi \cdot (x - x_0)$ for all $x \in A$. The *superdifferential* $\partial g(x_0)$ is the set of all supergradients.

Given a set E , the two superdifferentials $\partial\nu(E)$ and $\partial g(P(E))$ are related by the following simple lemma, that we report for later reference.

Lemma 3 *Given a measure game $\nu = g(P) : \Sigma \rightarrow \mathbb{R}$, for each set $E \in \Sigma$ a charge of the form $\chi \cdot P$ belongs to $\partial\nu(E)$ if and only if the vector $\chi \in \mathbb{R}^N$ belongs to $\partial g(P(E))$.*

Proof. The “if” part is trivial. As to the converse, suppose $\chi \cdot P \in \partial\nu(E)$. By definition, for all $A \in \Sigma$ we have:

$$g(P(A)) \leq g(P(E)) + \chi \cdot P(A) - \chi \cdot P(E) = g(P(E)) + \chi \cdot [P(A) - P(E)],$$

as desired. ■

We now present few elementary properties of the superdifferential $\partial\nu(E)$. It is easy to check that the set $\partial\nu(E)$ is convex and weak*-closed, and that the following properties hold:

- (i) $\partial\lambda\nu(E) = \lambda\partial\nu(E)$ for all $\lambda > 0$ and all $E \in \Sigma$.
- (ii) $\partial\nu_1(E) + \partial\nu_2(E) \subseteq \partial(\nu_1 + \nu_2)(E)$ for all $E \in \Sigma$ and all games ν_1 and ν_2 , with equality if at least one of the two games is in $ba(\Omega)$.

Given $E \in \Sigma$, consider the cone K_E defined by

$$K_E = \{m \in ba(\Omega) : m(G) \geq 0 \text{ and } m(F) \leq 0 \text{ for each } F \subseteq E^c \text{ and } G \subseteq E\}.$$

Clearly, $K_\Omega = ba(\Omega)^+$ and $-K_E = K_{E^c}$. Moreover, for all $m \in K_E$ and all $A \in \Sigma$ we have $m^+(A) = m(A \cap E)$ and $m^-(A) = -m(A \cap E^c)$, and so $\|m\| = 2m(E) - m(\Omega)$. The following result shows the importance of these cones for our analysis.

Proposition 4 *Let $\nu : \Sigma \rightarrow \mathbb{R}$ be a game superdifferentiable at E . Then, $\partial\nu(E) = \partial\nu(E) + K_{E^c}$ for each $E \in \Sigma$.*

Proof. The inclusion $\partial\nu(E) \subseteq \partial\nu(E) + K_{E^c}$ is obvious. As to the opposite inclusion, let $m \in \partial\nu(E) + K_{E^c}$. Then, for suitable $m_1 \in \partial\nu(E)$ and $m_2 \in K_{E^c}$, we have, for all $A \in \Sigma$,

$$\begin{aligned} \nu(E) + m(A) - m(E) &= \nu(E) + m_1(A) + m_2(A) - m_1(E) - m_2(E) \\ &\geq \nu(A) + m_2(A) - m_2(E) \\ &= \nu(A) + m_2(E^c \cap A) - m_2(E \cap A^c) \geq \nu(A), \end{aligned}$$

and so $m \in \partial\nu(E)$. ■

We now consider two key properties of superdifferentials, nonemptiness and the sum rule. Our first result shows that for the important class of exact games the set $\partial\nu(E)$ is nonempty for all $E \in \Sigma$.

Proposition 5 *If the game $\nu : \Sigma \rightarrow \mathbb{R}$ is exact, then $\partial\nu(E) \neq \emptyset$ for all $E \in \Sigma$. In particular, ν is exact if and only if $\partial\nu(E) \cap \text{core}(\nu) \neq \emptyset$ for all $E \in \Sigma$.*

Proof. Let ν be exact and let $E \in \Sigma$. By definition, there exists $m \in \text{core}(\nu)$ such that $m(E) = \nu(E)$. Since $m(A) \geq \nu(A)$ for all $A \in \Sigma$, it follows that $m \in \partial\nu(E)$, and so $\partial\nu(E) \cap \text{core}(\nu) \neq \emptyset$. Conversely, suppose that $\partial\nu(E) \cap \text{core}(\nu) \neq \emptyset$ for each $E \in \Sigma$. It is easy to check that $m \in \partial\nu(E) \cap \text{core}(\nu)$ implies $m(E) = \nu(E)$, and so ν is exact. ■

Since positive convex games are exact, they are superdifferentiable at all sets $E \in \Sigma$ by Proposition 5. The next result provides a simple extension of this result to real-valued convex games, thus providing a condition under which real-valued convex games are everywhere superdifferentiable.

Proposition 6 *A bounded game $\nu : \Sigma \rightarrow \mathbb{R}$ is convex if and only if $\partial\nu(E_1) \cap \partial\nu(E_2) \neq \emptyset$ for every pair $E_1 \subseteq E_2$.*

The next result shows that superdifferentials preserve sums.

Theorem 7 *Given any two convex and bounded games $\nu_1 : \Sigma \rightarrow \mathbb{R}$ and $\nu_2 : \Sigma \rightarrow \mathbb{R}$, we have*

$$\partial(\nu_1 + \nu_2)(E) = \partial\nu_1(E) + \partial\nu_2(E), \quad (2)$$

for all $E \in \Sigma$.

Since $\partial\lambda\nu(E) = \lambda\partial\nu(E)$ for all $\lambda > 0$ and all sets E , we conclude that, by Theorem 7, superdifferentials of bounded convex games preserve positive linear combinations. This fundamental property immediately implies the following interesting result, which shows that cores of bounded convex games are stable under summation.

Corollary 8 *Let $\nu_1 : \Sigma \rightarrow \mathbb{R}$ and $\nu_2 : \Sigma \rightarrow \mathbb{R}$ be any two convex and bounded games. If ν_1 and ν_2 have nonempty cores, then*

$$\text{core}(\nu_1 + \nu_2) = \text{core}(\nu_1) + \text{core}(\nu_2).$$

Proof. The nontrivial inclusion to prove is $\text{core}(\nu_1 + \nu_2) \subseteq \text{core}(\nu_1) + \text{core}(\nu_2)$. Let $m \in \text{core}(\nu_1 + \nu_2)$. Clearly, $m \in \partial(\nu_1 + \nu_2)(\emptyset)$ and $m(\Omega) = (\nu_1 + \nu_2)(\Omega)$. By Proposition 7, $m \in \partial\nu_1(\emptyset) + \partial\nu_2(\emptyset)$. Hence, there are $m_1 \in \partial\nu_1(\emptyset)$ and $m_2 \in \partial\nu_2(\emptyset)$ such that $m = m_1 + m_2$. By definition, $m_1 \geq \nu_1$ and $m_2 \geq \nu_2$. Hence, $m(\Omega) = (\nu_1 + \nu_2)(\Omega)$ implies $m_1(\Omega) = \nu_1(\Omega)$ and $m_2(\Omega) = \nu_2(\Omega)$. We conclude that $m_1 \in \text{core}(\nu_1)$ and $m_2 \in \text{core}(\nu_2)$, and so $m \in \text{core}(\nu_1) + \text{core}(\nu_2)$. ■

We close by considering measure games. In this case it is enough to study the existence of the standard superdifferential $\partial g(P(E))$ since, by Lemma 3, $\partial\nu(E)$ is nonempty whenever $\partial g(P(E))$ is nonempty.

Proposition 9 *Let $\nu = g(P) : \Sigma \rightarrow \mathbb{R}$ be a measure game. Then $\partial\nu(E) \neq \emptyset$ for all $E \in \Sigma$ if one of the following conditions holds:*

- (i) $g : R(P) \rightarrow \mathbb{R}$ can be extended as a concave function on some open convex set U containing $R(P)$.
- (ii) ν is superadditive and $g : R(P) \rightarrow \mathbb{R}$ is such that $g(\alpha P(E)) = \alpha g(P(E))$ for each $\alpha \in (0, 1)$ and each $E \in \Sigma$.

Condition (i) is the familiar sufficient condition from Convex Analysis, while condition (ii) is important in cooperative game theory, where the TU games that satisfy condition (ii) are called market games. They play an important role in the study of exchange economies (see Hart and Neyman, 1988).

4 Cores and Derivatives

The derivative for games introduced in Definition 1 was used by Epstein and Marinacci (2000) to study the cores of some TU games. In particular, they study a class of important economic games that have singleton cores and, loosely speaking, they show that the singleton actually consists of the derivative of the game. Since for real-valued functions the derivative can be viewed as a singleton superdifferential, it is natural to wonder whether the superdifferentials for games that we introduced are related to the derivatives of Definition 1, and, more importantly, whether they can be used to characterize cores that are not necessarily singleton. In this section we address these natural queries.

Interestingly, as in Epstein and Marinacci (2000), also in this work linear sets play a key role. Our first result provides a subcalculus characterization of the core based on linear sets.

Theorem 10 *Consider the following conditions:*

- (i) $E \in \mathcal{A}$.
- (ii) $\text{core}(\nu) = \partial\nu(E) \cap \partial\nu(E^c)$.
- (iii) $\partial\nu(E) \cap \partial\nu(E^c) \neq \emptyset$.

We have that (i) implies (ii), while the three conditions are equivalent whenever $\text{core}(\nu) \neq \emptyset$.

In other words, $\text{core}(\nu) = \partial\nu(E) \cap \partial\nu(E^c)$ when E is linear, regardless of whether or not $\text{core}(\nu)$ is nonempty. However, if $\text{core}(\nu)$ is nonempty, the three conditions are equivalent.

Proof. We first prove that (i) implies (ii). Let $E \in \mathcal{A}$. It is easy to see that $\text{core}(\nu) \subseteq \partial\nu(E) \cap \partial\nu(E^c)$. In fact, for each $m \in \text{core}(\nu)$ it holds that $m \geq \nu$, $m(E) = \nu(E)$, and $m(E^c) = \nu(E^c)$. We now prove the converse inclusion, that is, $\partial\nu(E) \cap \partial\nu(E^c) \subseteq \text{core}(\nu)$. Let $m \in \partial\nu(E) \cap \partial\nu(E^c)$. Since

$$0 = \nu(\emptyset) \leq \nu(E) - m(E),$$

we have $m(E) \leq \nu(E)$. Moreover, since

$$\nu(\Omega) = \nu(E \cup E^c) \leq \nu(E) + m(E^c),$$

we have $m(E^c) \geq \nu(E^c)$. By taking E^c in place of E , a similar argument shows that $m(E) \geq \nu(E)$ and $m(E^c) \leq \nu(E^c)$, and we conclude that $m(E) = \nu(E)$ and $m(E^c) = \nu(E^c)$. Finally, each set $B \in \Sigma$ can be written as $B = E \cup F - G$, with $F \cap E = \emptyset$ and $G \subseteq E$. Hence,

$$\begin{aligned}\nu(B) &= \nu(E \cup F - G) \leq \nu(E) + m(F) - m(G) \\ &= m(E) + m(F) - m(G) = m(B),\end{aligned}$$

and so $m \in \text{core}(\nu)$.

Next we prove that (iii) implies (i) when $\text{core}(\nu) \neq \emptyset$. Let $m \in \partial\nu(E) \cap \partial\nu(E^c)$. As we have just seen, this implies that $m(E^c) \leq \nu(E^c)$ and that $\nu(\Omega) \leq \nu(E) + m(E^c)$. Since $\text{core}(\nu) \neq \emptyset$, we have $\nu(E) + \nu(E^c) \leq \nu(\Omega)$. Hence,

$$\nu(\Omega) \leq \nu(E) + m(E^c) \leq \nu(E) + \nu(E^c) \leq \nu(\Omega),$$

and we conclude that $E \in \mathcal{A}$. To complete the proof, observe that (ii) obviously implies (iii) when $\text{core}(\nu) \neq \emptyset$. ■

Having established a subcalculus characterization of the core, we now move to study the relations of supergradients with the derivatives of games.

Theorem 11 *Let $\nu : \Sigma \rightarrow \mathbb{R}$ be a game superdifferentiable and differentiable at E . Then*

$$\delta\nu(\cdot; E) \in \partial\nu(E) + K_E.$$

If, in addition, E is linear and $\text{core}(\nu) \neq \emptyset$, then

$$\delta\nu(\cdot; E) \in \partial\nu(E^c).$$

Proof. Suppose that E is a maximum set for ν , that is, $\nu(E) \geq \nu(A)$ for all $A \in \Sigma$. Then, $\delta\nu(\cdot; E) \in K_E$. In fact, we have:

$$\begin{aligned}0 &= \lim_{\lambda \rightarrow 0} \sum_{j=1}^{n_\lambda} |\nu(E \cup F^{j,\lambda} - G^{j,\lambda}) - \nu(E) - \delta\nu(F^{j,\lambda}; E) + \delta\nu(G^{j,\lambda}; E)| \\ &\geq \lim_{\lambda \rightarrow 0} \sum_{j=1}^{n_\lambda} \nu(E) - \nu(E \cup F^{j,\lambda} - G^{j,\lambda}) + \delta\nu(F^{j,\lambda}; E) - \delta\nu(G^{j,\lambda}; E) \\ &\geq \lim_{\lambda \rightarrow 0} \sum_{j=1}^{n_\lambda} \delta\nu(F^{j,\lambda}; E) - \delta\nu(G^{j,\lambda}; E) = \delta\nu(F; E) - \delta\nu(G; E),\end{aligned}$$

and so $\delta\nu(G; E) \geq \delta\nu(F; E)$ for each $F \subseteq E^c$ and $G \subseteq E$. In particular, $\delta\nu(G; E) \geq \delta\nu(\emptyset; E) \geq \delta\nu(F; E)$, and we conclude that $\delta\nu(\cdot; E) \in K_E$. Let $m \in \partial\nu(E)$. By definition, $\nu - m$ is a game with maximum at E . Hence, by what we just proved, $\delta(\nu - m)(\cdot; E) \in K_E$, which implies $\delta\nu(\cdot; E) \in m(\cdot) + K_E$, and so $\delta\nu(\cdot; E) \in \partial\nu(E) + K_E$.

Suppose that $E \in \mathcal{A}$ and that $\text{core}(\nu) \neq \emptyset$. Let $m \in \text{core}(\nu)$. The game $\nu - m$ is a game with maximum at E . Hence, $\delta(\nu - m)(\cdot; E) \in K_E$, which, by Theorem 10 and Proposition 4, implies

$$\begin{aligned} \delta\nu(\cdot; E) &\in \text{core}(\nu) + K_E = \partial\nu(E) \cap \partial\nu(E^c) + K_E \\ &\subseteq \partial\nu(E^c) + K_E = \partial\nu(E^c), \end{aligned}$$

as desired. ■

In the last theorem we saw that $\delta\nu(\cdot; E) \in \partial\nu(E^c)$ when E is linear and $\text{core}(\nu) \neq \emptyset$. This raises the question of when $\delta\nu(\cdot; E) \in \partial\nu(E)$, something that in standard subcalculus happens in many important cases.

Theorem 12 *Let $\nu : \Sigma \rightarrow \mathbb{R}$ be a game differentiable at a linear set A . If $\text{core}(\nu) \neq \emptyset$, then $\partial\nu(A) \neq \emptyset$ and the following conditions are equivalent:*

- (i) $\delta\nu(\cdot; A) \in \partial\nu(A)$.
- (ii) $\text{core}(\nu) = \{\delta\nu(\cdot; A)\}$.
- (iii) $\delta\nu(\cdot; A) \in \text{core}(\nu)$.
- (iv) $\delta\nu(A; A) = \nu(A)$ and $\delta\nu(A^c; A) = \nu(A^c)$.

Moreover, if (i) and (iv) hold for some linear set A , then $\text{core}(\nu)$ is nonempty and coincides with the singleton $\{\delta\nu(\cdot; A)\}$.

Proof. Since A is linear, by Theorem 10 $\text{core}(\nu) \subseteq \partial\nu(A)$ and so $\text{core}(\nu) \neq \emptyset$ implies $\partial\nu(A) \neq \emptyset$. Having established that ν is superdifferentiable at A , we can prove that (i) implies (iv). Let $m \in \text{core}(\nu)$. By Theorem 10, $m \in \partial\nu(A^c)$, and so $\nu(A) \leq \nu(A^c) + m(A) - m(A^c)$. Moreover, $\delta\nu(\cdot; A) \in \partial\nu(A)$ implies $\nu(A^c) \leq \nu(A) + \delta\nu(A^c; A) - \delta\nu(A; A)$. Adding up, we get $\delta\nu(A; A) - m(A) \leq \delta\nu(A^c; A) - m(A^c)$. On the other hand, since $\nu - m$ has

a maximum at A , by what we proved in the proof of Theorem 11 we have $\delta(\nu - m)(\cdot; A) \in K_A$, and so

$$\delta\nu(A; A) - m(A) \geq 0 \geq \delta\nu(A^c; A) - m(A^c).$$

All this implies that $\delta\nu(A; A) - m(A) = \delta\nu(A^c; A) - m(A^c) = 0$. Since $A \in \mathcal{A}$ and $m \in \text{core}(\nu)$, we conclude that $\delta\nu(A; A) = \nu(A)$ and $\delta\nu(A^c; A) = \nu(A^c)$.

We now show that (iv) implies (ii). Let $m \in \text{core}(\nu)$. Since $\delta(\nu - m)(\cdot; A) \in K_A$, then for each $G \subseteq A$ it holds that $\delta\nu(G; A) \geq m(G)$. Along with $\delta\nu(A; A) = \nu(A) = m(A)$, this implies that $m(G) = \delta\nu(G; A)$ for each $G \subseteq A$. A similar argument shows that $m(F) = \delta\nu(F; A)$ for each $F \subseteq A^c$, and so $m = \delta\nu(\cdot; A)$, which proves that $\text{core}(\nu) = \{\delta\nu(\cdot; A)\}$.

Since (ii) trivially implies (iii), it remains to prove that (iii) implies (i). Since $\text{core}(\nu) \neq \emptyset$ and $A \in \mathcal{A}$, by Theorem 10 $\delta\nu(\cdot; A) \in \text{core}(\nu) = \partial\nu(A) \cap \partial\nu(A^c) \subseteq \partial\nu(A)$.

Suppose that (i) and (iv) hold for some $A \in \mathcal{A}$. By (iv), $\nu(\Omega) = \delta\nu(\Omega; A)$ and together (i) and (iv) imply that, for all $E \in \Sigma$,

$$\nu(E) \leq \delta\nu(E; A) + \nu(A) - \delta\nu(A; A) = \delta\nu(E; A).$$

Hence, $\delta\nu(\cdot; A) \in \text{core}(\nu)$ and so, by what we proved above, $\text{core}(\nu) = \{\delta\nu(\cdot; A)\}$. ■

Example. Let $\nu = g(P) : \Sigma \rightarrow \mathbb{R}$ be a measure game and suppose there is a linear set A such that g is differentiable and superdifferentiable at $P(A) \in \text{ri}(R(P))$. By Theorem 12, $\text{core}(\nu) \subseteq \{\nabla g(P(A)) \cdot P(\cdot)\}$. In fact, since $P(A) \in \text{ri}(R(P))$, by a well-known result of Convex Analysis for each $\chi \in \partial g(P(A))$ we have $[\nabla g(P(A)) - \chi] \cdot w = 0$ for all $w \in \text{span}(R(P))$. Hence, $\nabla g(P(A)) \cdot P(\cdot) = \chi \cdot P(\cdot)$ and so, by Lemma 3, $\nabla g(P(A)) \cdot P(\cdot) \in \partial g(P(A))$. Since in Epstein and Marinacci (2000) it is shown that $\delta\nu(\cdot; A) = \nabla g(P(A)) \cdot P(\cdot)$, we conclude that $\delta\nu(\cdot; A) \in \partial\nu(A)$ and so, by Theorem 12, $\text{core}(\nu) \subseteq \{\nabla g(P(A)) \cdot P(\cdot)\}$.

Example. Consider the production game $\nu = g(P) : \Sigma \rightarrow \mathbb{R}$ defined via the Cobb-Douglass function $g(x_1, \dots, x_N) = x_1^{\alpha_1} \cdots x_N^{\alpha_N}$, where each $\alpha_i \geq 0$ and $\sum_{i=1}^N \alpha_i = 1$. If $P(\Omega) \in \mathbb{R}_{++}^N$, then, by Theorem 12,

$$\text{core}(\nu) = \{g(P(\Omega))(\nabla g(P(A)) \cdot P(\cdot))\} = \left\{ g(P(\Omega)) \sum_{i=1}^N \alpha_i P_i(\cdot) \right\}. \quad (3)$$

For, given any $P(\Omega) \in \mathbb{R}_{++}^N$, the concave function g is differentiable and superdifferentiable at all points of $ri(R(P))$. In particular, it is easy to check that all sets A_t such that $P(A_t) = tP(\Omega)$ for some $t \in (0, 1)$ are linear and belong to $ri(R(P))$. Hence, by the argument of the previous example, $\delta\nu(\cdot; A_t) \in \partial\nu(A_t)$. Moreover, some simple algebra shows that $\delta\nu(A_t; A_t) = \nu(A_t)$ and $\delta\nu(A_t^c; A_t) = \nu(A_t^c)$. Hence, conditions (i) and (iv) of Theorem 12 are therefore satisfied and so, by Theorem 12, Eq. (3) holds.

5 Measure Games

Games relevant for economic applications have often the form of a measure game $g(P) : \Sigma \rightarrow \mathbb{R}$. In this section we study in more detail the structure of the superdifferentials and cores of this class of games.

The natural question for cores of measure games is how to relate the underlying vector charge P with the charges in the cores. We start by establishing a general result of this type for the important countably additive case. To this end, we introduce lower Lipschitz functions.

Definition 13 *A function $g : A \subseteq \mathbb{R}^N \rightarrow \mathbb{R}$ is (locally) lower Lipschitzian at $x_0 \in A$ if there is a neighborhood $B = B(x_0, \varepsilon)$ of x_0 and a constant $\gamma > 0$ such that, for all $x \in B \cap A$,*

$$[g(x) - g(x_0)]^- \leq \gamma \|x - x_0\|.$$

We denote by P^* the “average” measure $(1/N) \sum_{i=1}^N P_i$ and in the next statement $\|\cdot\|_\infty$ is the norm of the space $L^\infty(\Omega, \Sigma, P^*)$. In reading the result keep in mind that a function is lower semicontinuous at some point when it is there lower Lipschitzian.

Theorem 14 *Let $g(P) : \Sigma \rightarrow \mathbb{R}$ be a measure game and suppose that P is countably additive and that there is a linear set A (e.g., $A = \emptyset$) such that g is lower semicontinuous at $P(A)$ and $P(A^c)$. Then, for each $m \in \text{core}(\nu)$ there exists a Σ -measurable vector function $f = (f_1, \dots, f_N) : \Omega \rightarrow \mathbb{R}^N$ such that, for all $E \in \Sigma$,*

$$m(E) = \sum_{i=1}^N \int_E f_i dP_i. \quad (4)$$

Moreover, a Σ -measurable vector function $f : \Omega \rightarrow \mathbb{R}^N$ satisfies (4) if and only if

$$\sum_{i=1}^N \frac{dP_i}{dP^*} f_i = \frac{dm}{dP^*} \quad P^* \text{-a.e.} \quad (5)$$

If, in addition, g is lower Lipschitzian at 0 and $P(\Omega)$, then there exists $\gamma > 0$ such that $\|dm/dP^*\|_\infty \leq \gamma$ for all $m \in \text{core}(\nu)$.

Notice that Eq. (4) provides two important pieces of information on the charges belonging to $\text{core}(\nu)$: (i) they are all countably additive; (ii) they are all absolutely continuous w.r.t. P^* .

An especially interesting case in Theorem 14 is when to a given m in $\text{core}(\nu)$ corresponds a constant vector function $f : \Omega \rightarrow \mathbb{R}^N$, that is, when there exists a vector $(\alpha_1, \dots, \alpha_N) \in \mathbb{R}^N$ such that $f(\omega) = (\alpha_1, \dots, \alpha_N)$ for all $\omega \in \Omega$. In this case, m is a linear combination of the underlying vector charge P , a most convenient situation.

Because of their interest, we first give a name to the subset of $\text{core}(\nu)$ consisting of such linear combinations.

Definition 15 *The linear core of a measure game $\nu : g(P) : \Sigma \rightarrow \mathbb{R}$ is the subset $\mathcal{L}\text{core}(\nu)$ of $\text{core}(\nu)$ defined by*

$$\mathcal{L}\text{core}(\nu) = \text{core}(\nu) \cap \text{span}\{P_1, \dots, P_N\}.$$

Using Lemma 3 and Theorem 10, it is easy to characterize the linear core and to provide bounds for its dimension. All this makes use of linear sets, thus showing their importance for $\mathcal{L}\text{core}(\nu)$.

Proposition 16 *Given a measure game $\nu = g(P) : \Sigma \rightarrow \mathbb{R}$, it holds that*

$$\mathcal{L}\text{core}(\nu) = \{\chi \cdot P : \chi \in \partial g(P(A)) \cap \partial g(P(A^c))\} \quad (6)$$

for each linear set A . Moreover,

$$\dim(\mathcal{L}\text{core}(\nu)) \leq \dim(R(P)) - \dim(\text{span}\{P(A) : A \in \mathcal{A}\}) \leq N - 1. \quad (7)$$

Remark. This result holds for any vector charge P , not necessarily strongly continuous. In this more general setting $R(P)$ may not be convex and, therefore, $\dim(R(P))$ has to be replaced with $\dim(\text{span}(R(P)))$.

Example. By Eq. (7), if $\nu = g(P) : \Sigma \rightarrow \mathbb{R}$ is a scalar measure game, then $\mathcal{L}core(\nu)$, when nonempty, is a singleton with $\chi = g(P(\Omega)) / P(\Omega)$. In particular, suppose that g is positive and continuous at $P(\Omega)$ and that P is countably additive. Then, it can be checked that each $m \in core(\nu)$ is countably additive and non-atomic. A simple application of the Lyapunov Theorem shows that $core(\nu) \neq \emptyset$ if and only if $(g(P(\Omega)) / P(\Omega)) x \geq g(x)$ for all $x \in [0, P(\Omega)]$. For, let $m \in core(\nu)$ and consider the vector measure (m, P) . By the Lyapunov Theorem, given any $x \in [0, P(\Omega)]$ there is E such that $(P(E), m(E)) = (x, (g(P(\Omega)) / P(\Omega)) x)$. Hence, $(g(P(\Omega)) / P(\Omega)) x = m(E) \geq g(P(E)) = g(x)$. All this implies that in this case $\mathcal{L}core(\nu) \neq \emptyset$ if and only if $core(\nu) \neq \emptyset$.

6 Linear Games

In the last section we introduced the linear core, the subset of the core of a measure game $g(P) : \Sigma \rightarrow \mathbb{R}$ that consists of linear combinations of the underlying vector charge P . This part of the core is especially interesting because of its simple form and analytical tractability, and the games whose core and linear core coincide stand out among games in terms of simplicity and tractability. This section is devoted to the study of these games, that we call linear.

Definition 17 *A measure game $\nu = g(P) : \Sigma \rightarrow \mathbb{R}$ is called linear if $core(\nu) = \mathcal{L}core(\nu)$, that is, if $core(\nu) \subseteq span\{P_1, \dots, P_N\}$.*

To provide a characterization of linear games we first state a linearity theorem for vector measures that should be of independent interest. The following important class of sets will play a key role.

Definition 18 *A set $A \in \Sigma$ is radial if there is a set $E \in \Sigma$ such that, for some $t \in (0, 1)$,*

$$P(A) = tP(E) + (1 - t)P(E^c).$$

By the Lyapunov Theorem, radial sets form a significant subset of $R(P)$ and they include the sets called *diagonal* by Epstein and Marinacci (2000), that is, the sets $A \in \Sigma$ such that $P(A) = tP(\Omega)$ for some $t \in (0, 1)$. The next result provides a useful characterization of radial sets in terms of the relative interior of $R(P)$. It is based on the important property of the range $R(P)$ of

having the point $2^{-1}P(\Omega)$ as a center of symmetry, that is, $2(2^{-1}P(\Omega)) - x \in R(P)$ for all $x \in R(P)$.³ This “pivotal” feature of $2^{-1}P(\Omega)$ is key in the next result.

Proposition 19 *Let Σ be a σ -algebra of subsets, and $P = (P_1, \dots, P_N) : \Sigma \rightarrow \mathbb{R}^N$ a vector charge with each P_i strongly continuous. Then, a set $E \in \Sigma$ is radial if and only if $P(E)$ belongs to the relative interior of $R(P)$.*

We can now state and prove the announced linearity theorem.

Theorem 20 *Let Σ be a σ -algebra of subsets, $P = (P_1, \dots, P_N) : \Sigma \rightarrow \mathbb{R}_+^N$ a positive vector charge with each P_i strongly continuous and suppose $m : \Sigma \rightarrow \mathbb{R}$ is either a signed measure in $ca(\Omega)$ or a strongly continuous charge in $ba(\Omega)$. If there exists a radial set A such that, for all $E \in \Sigma$,*

$$P(E) = P(A) \implies m(E) = m(A), \quad (8)$$

then

$$m \in \text{span} \{P_1, \dots, P_n\}. \quad (9)$$

If, in addition, it holds that

$$P(E) \geq P(A) \implies m(E) \geq m(A), \quad (10)$$

then

$$m \in \text{cone} \{P_1, \dots, P_n\}.$$

Finally, the coefficients for which (9) holds are unique if and only if $R(P)$ is full dimensional.

It is important to note the two key features of this result: (i) the existence of just a single radial set A is required; (ii) no assumption, besides either countable additivity or strong continuity, is made on m . Theorem 20 is the N -dimensional generalization of a uniqueness result of Marinacci (1998) which holds for positive scalar measures P and m . In fact, in the scalar case a set A is radial if and only if $0 < P(A) < P(\Omega)$. Therefore, if there exists a set $A \in \Sigma$ with $0 < P(A) < P(\Omega)$ and such that

$$P(E) = P(A) \implies m(E) = m(A)$$

³See, e.g., Bolker (1969), who studies in detail the geometry of $R(P)$.

whenever $E \in \Sigma$, then $m(E) = \frac{m(\Omega)}{P(\Omega)}P(E)$ by Theorem 20. When m is positive, this is the uniqueness result of Marinacci (1998). In that paper, however, uniqueness is also proved for lambda systems, while here we only consider σ -algebras.

Though in this paper we focus on TU games, Theorem 20 can be also interpreted in a social choice context if we assume that m and each P_i are probability measures representing beliefs. For instance, consider diagonal sets, that in this setting can be viewed as events over which agents have unanimous beliefs, say $P_i(A) = \alpha \in (0, 1)$ for each $i = 1, \dots, N$. By Theorem 20, linear aggregation occurs whenever the aggregator m preserves the agents' unanimous beliefs on some event A , a condition much weaker than the Paretian conditions used in Bayesian aggregation results (cf. Fishburn, 1984 and Mongin, 1995).

Two final remarks:

(i) Two-dimensional vector probabilities $(P_1, P_2) : \Sigma \rightarrow [0, 1]$ have full dimensional range provided that just $P_1 \neq P_2$. In the N -dimensional case, $(P_1, \dots, P_N) : \Sigma \rightarrow \mathbb{R}_+^N$ has full range when, for example, the measures P_i are mutually singular.

(ii) Theorem 20 holds even when m is a vector measure $m : \Sigma \rightarrow \mathbb{R}^M$. In this case, there exists a $M \times N$ matrix of coefficients A such that $m = AP$.

6.1 Characterizing Linear Games

Using Theorem 10 and the just established Theorem 20, we can now provide a simple condition under which a measure game is linear.

Theorem 21 *Let $\nu = g(P) : \Sigma \rightarrow \mathbb{R}$ be a measure game and suppose one of the following holds:*

(i) *P is countably additive and there is a linear set A_* (e.g., $A_* = \emptyset$) such that g is lower semicontinuous at $P(A_*)$ and $P(A_*^c)$.*

(ii) *g is lower Lipschitzian at 0 and $P(\Omega)$.*

Then, if there exists a linear and radial set, the game ν is linear and

$$\text{core}(\nu) = \{\chi \cdot P : \chi \in \partial g(P(A)) \cap \partial g(P(A^c))\} \quad (11)$$

for each linear sets $A \in \Sigma$. The vectors χ are univocally determined if and only if $R(P)$ is full dimensional, while if g is monotone on $R(P)$, then χ

can be assumed to be non-negative, i.e., $\chi_i \geq 0$ for all $1 \leq i \leq N$. Finally, if ν is exact the converse holds, that is, a linear and exact measure game has linear and radial sets.

Remarks. (i) In Eq. (11) it may well happen that

$$\text{core}(\nu) = \{\chi \cdot P : \chi \in \partial g(P(A)) \cap \partial g(P(A^c))\} = \emptyset.$$

(ii) The converse does not hold if ν is not exact. In fact, consider the following scalar measure game:

$$\nu(E) = \begin{cases} P(E) & \text{if } P(E) < \frac{1}{2} \\ P(E)^2 & \text{if } P(E) \geq \frac{1}{2} \end{cases} \quad (12)$$

with $P(\Omega) = 1$. It is easy to check that $\text{core}(\nu) = \{P\}$. However, there are no radial sets that are linear, i.e., there are no sets A such that $P(A) \in (0, 1)$ and $\nu(A) + \nu(A^c) = 1$.

Proof. Let A be linear and radial. By Theorem 10, $\text{core}(\nu) = \partial\nu(A) \cap \partial\nu(A^c)$. Suppose $\text{core}(\nu) \neq \emptyset$. For all $m \in \text{core}(\nu)$ and for all $E \in \Sigma$ we have:

$$\begin{aligned} g(P(E)) &\leq g(P(A)) + m(E) - m(A), \\ g(P(E^c)) &\leq g(P(A^c)) + m(E^c) - m(A^c). \end{aligned}$$

Hence, $P(E) = P(A)$ implies $m(E) = m(A)$, and so Eq. (8) of Theorem 20 holds. On the other hand, if (i) holds, then, by Lemma 35, all charges in $\text{core}(\nu)$ are countably additive, while if (ii) holds, then, by Lemma 34, all such charges are strongly continuous. In both cases we can apply Theorem 20, and we conclude that $m \in \text{span}\{P_1, \dots, P_N\}$. The game is linear and, by Proposition 16, $\chi \in \partial g(P(A)) \cap \partial g(P(A^c))$.

Now, let A be any linear set. By Proposition 16,

$$\text{core}(\nu) = \mathcal{L}\text{core}(\nu) = \{\chi \cdot P : \chi \in \partial g(P(A)) \cap \partial g(P(A^c))\},$$

which proves Eq. (11).

If g is monotone, then for all $m \in \text{core}(\nu)$ and for all $E \in \Sigma$ we have:

$$m(E) - m(A) \geq g(P(E)) - g(P(A)) \geq 0$$

whenever $P(E) \geq P(A)$. Hence, Eq. (10) of Theorem 20 holds and, therefore, the vector χ can be chosen to be non-negative.

The only nontrivial part it remains to prove is that exact linear games have radial and linear sets. Given $t \in (0, 1)$, let A be the diagonal set such that $P(A) = tP(\Omega)$. Given $m \in \text{core}(\nu)$, for a suitable $\chi \in \mathbb{R}^N$ we have $m(A) = \chi \cdot P(A) = t\chi \cdot P(\Omega) = t\nu(\Omega)$. Hence, $m(A) = t\nu(\Omega)$ for all $m \in \text{core}(\nu)$ and so, by exactness, $\nu(A) = t\nu(\Omega)$. A similar argument shows that $\nu(A^c) = (1 - t)\nu(\Omega)$, and we conclude that A is linear.

Finally, when $\text{core}(\nu) = \emptyset$, by Lemma 3 and Theorem 10 we have

$$\{\chi \cdot P : \chi \in \partial g(P(A)) \cap \partial g(P(A^c))\} \subseteq \text{core}(\nu)$$

for each linear set $A \in \Sigma$. ■

Conditions (i) and (ii) of Theorem 21 are both very mild requirements. In particular, condition (i) is more demanding on P , which is required to be countable additivity rather than just finite additive, but less on g , which is only required to be lower semicontinuous rather than lower Lipschitzian.

As to the existence of linear and radial sets, measure games $g(P)$ that are relevant for economic applications typically feature some homogeneity conditions of the function $g : R(P) \rightarrow \mathbb{R}$, and these conditions guarantee the existence of many linear and radial sets for the measure game $g(P)$.

For instance, say that the measure game $\nu = g(P) : \Sigma \rightarrow \mathbb{R}$ is *radially concave at E* if, for all $t \in (0, 1)$,

$$g(tP(E) + (1 - t)P(E^c)) \geq tg(P(E)) + (1 - t)g(P(E^c)). \quad (13)$$

Obviously, ν is radially concave at E if and only if it is radially concave at E^c , and ν is radially concave at all sets E in Σ when g is concave.

Definition 22 *A measure game $\nu = g(P) : \Sigma \rightarrow \mathbb{R}$ is called radially concave if there is some linear set A such that ν is radially concave at A .*

For example, since Ω is a linear set, ν is radially concave if, for all $t \in (0, 1)$,

$$g(tP(\Omega)) \geq tg(P(\Omega)),$$

a very mild homogeneity requirement. Another simple case in which ν is radially concave is when the set A such that $P(A) = 2^{-1}P(\Omega)$ is linear. In this case Eq. (13) is trivially satisfied.

Radial concavity is a weak condition satisfied by many economic TU games. For example, the measure games whose functions $g : R(P) \rightarrow \mathbb{R}$ are concave or homogeneous of degree one are radially concave, as well as the measure games that have a function $g : R(P) \rightarrow \mathbb{R}$ homogeneous of degree $k < 1$, provided $g(P(\Omega)) \geq 0$. In particular, market games are radially concave, as their function g is homogeneous of degree one.

Radially concave games that have nonempty cores admit many radial and linear sets, and, consequently, by Theorem 21, they are linear, as stated in the next Corollary.

Corollary 23 *Let $\nu = g(P) : \Sigma \rightarrow \mathbb{R}$ be a radially concave measure game and suppose one of conditions (i) and (ii) of Theorem 21 holds. Then, the game ν is linear and, for each linear set A ,*

$$\begin{aligned} \text{core}(\nu) &= \{\chi \cdot P : \chi \in \partial g(P(A)) \cap \partial g(P(A^c))\} \\ &= \{\chi \cdot P : \chi \in \partial g(2^{-1}P(\Omega))\}. \end{aligned}$$

Remark. Interestingly, here $\text{core}(\nu)$ is determined by the superdifferential of g at $2^{-1}P(\Omega)$, the center of symmetry of $R(P)$.

Example. Let $g : \mathbb{R}_+^N \rightarrow \mathbb{R}$ be a concave and positive homogeneous function and assume $P(\Omega) \in \mathbb{R}_{++}^N$. Consider the following two broad classes of functions:

$$\begin{aligned} g_1(x) &= g(x) + h_1(x), \\ g_2(x) &= g(x) h_2(x), \end{aligned}$$

for all $x \in \mathbb{R}_+^N$. If $h_1(tP(\Omega)) = 0$ and $h_2(tP(\Omega)) = 1$ for all $t \geq 0$, then the games $g_1(P)$ and $g_2(P)$ are radially concave. In view of Corollary 23, it is easy to provide conditions under which the cores of these measure games are nonempty. For instance, for the first class it suffices that $\partial h_1(2^{-1}P(\Omega)) \neq \emptyset$, while for the other class it is enough to require that $h_2(x) \in [0, 1]$ for all $x \in \mathbb{R}_+^N$.

In Theorem 21 and in Corollary 23 we only assumed that the real-valued function g was defined on the range $R(P)$. In applications, however, it is often the case that the function g defining the measure game is defined on an open convex subset G containing $R(P)$, for example \mathbb{R}^N itself. In this case, we have two superdifferentials, the one of g restricted to $R(P)$, i.e.

$\partial g|_{R(P)}(x)$, and the one that g has relative to the open convex subset G , i.e. $\partial g(x)$. Naturally, $\partial g|_{R(P)}(x)$ is the superdifferential relevant for Theorem 21 and Corollary 23. On the other hand, the superdifferential $\partial g(x)$ may be easier to compute, especially when g is defined on \mathbb{R}^N .

The next result can therefore be useful, as it shows that it is possible to use directly $\partial g(P(A))$ when g is concave and A radial.

Proposition 24 *Let $P = (P_1, \dots, P_N) : \Sigma \rightarrow \mathbb{R}$ be a strongly continuous vector charge and let $g : G \rightarrow \mathbb{R}$ be a concave function, where G is an open convex set containing $R(P)$. For the measure game $\nu = g(P) : \Sigma \rightarrow \mathbb{R}$, it holds that*

$$\begin{aligned} \text{core}(\nu) &= \{\chi \cdot P : \chi \in \partial g(P(A)) \cap \partial g(P(A^c))\} \\ &= \{\chi \cdot P : \chi \in \partial g(2^{-1}P(\Omega))\}. \end{aligned}$$

for each linear and radial set A .

Example (Generalized Linear Production Games). Let $a : T \rightarrow \mathbb{R}^N$ be a continuous map, where T is a compact metric space, and define a function $g : \mathbb{R}^N \rightarrow \mathbb{R}$ by $g(x) = \min_{t \in T} a(t) \cdot x$ for all $x \in \mathbb{R}^N$. Consider the measure game $\nu = g(P)$, which we call a *generalized linear production game*. When T is a finite set and $a(t) \equiv a^t \in \mathbb{R}_+^N$, we have the linear production games of Owen (1975) and Billera and Raanan (1981). Since the function g is concave on \mathbb{R}^N , by a standard result in Convex Analysis (see, e.g., Hiriart-Urruty and Lemarechal, 1993), we have

$$\partial g(x) = \text{co}(a(t) : t \in I(x)),$$

where $I(x) = \{t : a(t) \cdot x = g(x)\}$. Consider a diagonal set A with $P(A) = \alpha P(\Omega)$ for some $\alpha \in (0, 1)$. Simple algebra shows that

$$I(P(A)) = I(P(\Omega)) = \{t : a(t) \cdot P(\Omega) = g(P(\Omega))\}.$$

Since each diagonal set is linear, by Proposition 24,

$$\text{core}(\nu) = \{\chi \cdot P : \chi \in \text{co}(a(t) : a(t) \cdot P(\Omega) = \nu(\Omega))\}.$$

This includes Corollary 2.7 of Billera and Raanan (1981), which therefore follows from Proposition 24 using some standard Convex Analysis.

6.2 Differentiability

Proposition 16 characterized the linear core of a measure game $\nu = g(P) : \Sigma \rightarrow \mathbb{R}$ through the superdifferentials of the function $g : R(P) \rightarrow \mathbb{R}$ and Theorem 21 provided a simple condition under which the entire core can be characterized in this way. In view of standard subcalculus and of Theorem 12, it is natural to wonder what happens when some differentiability is assumed on g , in particular, whether the core shrinks to a singleton.

Proposition 25 *Let $\nu = g(P) : \Sigma \rightarrow \mathbb{R}$ be a measure game and suppose one of conditions (i) and (ii) of Theorem 21 holds. If there is a linear and radial set A such that g is differentiable at $P(A)$, then*

$$\text{core}(\nu) = \emptyset \quad \text{or} \quad \text{core}(\nu) = \{\nabla g(P(A)) \cdot P(\cdot)\}.$$

If, in addition, g is differentiable and superdifferentiable at both $P(A)$ and $P(A^c)$, then $\text{core}(\nu) \neq \emptyset$ if and only if $\nabla g(P(A)) = \nabla g(P(A^c))$.

Differentiability has therefore a remarkably strong impact on the core: even just assuming that g is differentiable at $P(A)$ forces the core to be at most a singleton.

Example. Let $\nu = g(P) : \Sigma \rightarrow \mathbb{R}$ be a market game, that is, ν is superadditive and g is homogeneous of degree one. If g is differentiable at $P(\Omega)$, then $\text{core}(\nu) = \{\nabla g(P(\Omega)) \cdot P(\cdot)\}$. In fact, by Proposition 9, $\partial g(P(E)) \neq \emptyset$ for all $E \in \Sigma$. Moreover, all diagonal sets are linear and g is differentiable at all them because it is differentiable at $P(\Omega)$. In particular, $\nabla g(P(A)) = \nabla g(P(\Omega))$ for all diagonal sets. Hence, by Proposition 25, $\text{core}(\nu) = \{\nabla g(P(\Omega)) \cdot P(\cdot)\} = \{\nabla g(P(\Omega)) \cdot P(\cdot)\}$. This result is essentially due to Aumann and Shapley (1974) and plays a key role in their analysis of exchange economies. It therefore follows from Proposition 25.

Example. Let $\nu = g(P) : \Sigma \rightarrow \mathbb{R}$ be a measure game defined as follows:

$$g(P(E)) = \begin{cases} 1 - \frac{2(1-P_1(E))^3}{(1-P_1(E))^2 + (1-P_2(E))^2} & \text{if } (P_1(E), P_2(E)) \neq (1, 1) \\ 0 & \text{if } P_1(E) = P_2(E) = 1 \end{cases}$$

All diagonal sets of this game are linear and g is differentiable at all them. Hence, by Proposition 25, $\text{core}(\nu) \subseteq \{2P_1 - P_2\}$. In fact, some algebra shows that $\text{core}(\nu) = \{2P_1 - P_2\}$.

Unlike Proposition 25, the next result does not require A to be radial, at the cost of a stronger assumption on the function g .

Corollary 26 *Let $\nu = g(P) : \Sigma \rightarrow \mathbb{R}$ be a measure game and suppose one of conditions (i) and (ii) of Theorem 21 holds. If there is a linear set A such that ν is radially concave at A and g is differentiable on some neighborhood U of $P(A)$, then*

$$\text{core}(\nu) = \emptyset \quad \text{or} \quad \text{core}(\nu) = \{\nabla g(P(A)) \cdot P(\cdot)\}.$$

Remark. If $g : R(P) \rightarrow \mathbb{R}$ is concave and differentiable at $P(A)$, then the corollary holds.⁴

Proof. Since ν is radially concave at A , it is easy to check that all sets A_α such that $P(A_\alpha) = \alpha P(A) + (1 - \alpha) P(A^c)$ are linear (see the proof of Corollary 23). For α small enough, $P(A_\alpha) \in U$ and so g is differentiable at $P(A_\alpha)$. Since A_α is a radial set, a simple application of Proposition 25 proves the result. ■

7 Finite-Dimensional Cores

Let $sc(\Omega)$ be the vector subspace spanned by the strongly continuous charges of $ba(\Omega)$. The key feature of linear measure games is that their cores are finite-dimensional subsets of $sc(\Omega)$. As in exact games there is a tight connection between the core and the game, one may wonder whether there exists a characterization of general exact games featuring finite-dimensional cores in $sc(\Omega)$.

In this section we provide such a characterization: Theorem 31 shows that an exact game has a finite-dimensional core in $sc(\Omega)$ if and only if it is a generalized linear production game. This is a class of measure games introduced in the previous section and whose associated function g has the very simple form $g(x) = \min_{t \in T} a(t) \cdot x$, where T is a compact metric space and $a : T \rightarrow \mathbb{R}^N$ is a continuous map.

The characterization is therefore sharp: exact generalized linear production games are the only exact games having finite-dimensional cores in $sc(\Omega)$. As Hart and Neyman (1988) observe on p. 32, most relevant economic TU

⁴Recall that g can be differentiable at $P(A)$ only if g is defined (or can be extended) on a suitable open subset of $P(A)$.

games are linear. In view of Theorem 31, we can say that all such games, when exact, have to be generalized linear production games.

To prove this characterization we have to introduce some notions, that may be of independent interest. We denote by Γ_α a (possibly finite) w^* -compact subset of $ba(\Omega)$ such that $m(\Omega) = \alpha$ for each $m \in \Gamma_\alpha$.

Definition 27 *The closure under majorization (m -closure, for short) $\tilde{\Gamma}_\alpha$ of a set Γ_α is defined as follows: $m \in \tilde{\Gamma}_\alpha$ if $m(\Omega) = \alpha$ and if for each $E \in \Sigma$ there is $m' \in \Gamma_\alpha$ such that $m(E) \geq m'(E)$.*

We say that a set Γ_α is m -closed if $\tilde{\Gamma}_\alpha = \Gamma_\alpha$. The next lemma contains a few properties of the m -closure.

Lemma 28 *For each set Γ_α we have:*

$$(i) \ w^* - cl(co(\Gamma_\alpha)) \subseteq \tilde{\Gamma}_\alpha \text{ and } \widetilde{(\tilde{\Gamma}_\alpha)} = \tilde{\Gamma}_\alpha.$$

(ii) $\tilde{\Gamma}_\alpha$ is w^* -compact and convex.

(iii) *The game*

$$\nu(\cdot) = \min_{m \in \tilde{\Gamma}_\alpha} m(\cdot) = \min_{m \in \Gamma_\alpha} m(\cdot)$$

is the unique exact game such that $core(\nu) = \tilde{\Gamma}_\alpha$.

An interesting consequence of Lemma 28 is that a w^* -compact and convex set is the core of a game if and only if it is m -closed. In fact, cores are clearly m -closed. On the other hand, by point (iii) of Lemma 28, a w^* -compact, convex, and m -closed set $\tilde{\Gamma}_\alpha$ is the core of the game $\min_{m \in \tilde{\Gamma}_\alpha} m(\cdot)$. Another noteworthy consequence of point (iii) is the existence of a one-to-one correspondence between exact games and the sets $\tilde{\Gamma}_\alpha$.

Let $\pi : \mathbb{R}^N \rightarrow span(R(P))$ be the orthogonal projection on $span(R(P))$. Define the map $R : span\{P_1, \dots, P_N\} \rightarrow span(R(P))$ between our two key vector spaces as follows: $R(\chi \cdot P) = \pi(\chi)$ for all $\chi \in \mathbb{R}^N$. This means that, for all $m \in span\{P_1, \dots, P_N\}$, we have $m = R(m) \cdot P$.

By the next result, R is a “canonical” isomorphism between the two spaces.

Lemma 29 *The map $R : span\{P_1, \dots, P_N\} \rightarrow span(R(P))$ is a linear and w^* -continuous isomorphism.*

Using the canonical isomorphism R , we get the following important property of the m -closure.

Lemma 30 *Let $P = (P_1, \dots, P_N) : \Sigma \rightarrow \mathbb{R}$ be a strongly continuous vector charge. Then $\Gamma_\alpha \subseteq \text{span} \{P_1, \dots, P_N\}$ implies $\tilde{\Gamma}_\alpha = \text{co}(\Gamma_\alpha)$.*

The next example shows that Lemma 30 may fail if P is not strongly continuous.

Example. Let $\Omega = \{\omega_1, \omega_2, \omega_3\}$ and $\Sigma = 2^\Omega$. Consider the compact and convex set of probability charges

$$\Gamma_1 = \left\{ m : \Sigma \rightarrow [0, 1] : \sum_{i=1}^3 m(\omega_i) = 1 \text{ and } m(\omega_1) \leq m(\omega_2) \right\}.$$

It is easy to check that the charge $m^* = (1/2, 0, 1/2)$ belongs to $\tilde{\Gamma}_1$, but it does not belong to $\text{co}(\Gamma_1)$. In particular, this implies that Γ_1 cannot be the core of any game.

We can now state and prove the announced characterization, which shows that an exact game has a finite-dimensional core in $sc(\Omega)$ if and only if it is a generalized linear production game.

Theorem 31 *Let $P = (P_1, \dots, P_N) : \Sigma \rightarrow \mathbb{R}_+^N$ be a strongly continuous vector charge. Given a game $\nu : \Sigma \rightarrow \mathbb{R}$, consider the following conditions:*

- (i) ν is exact and $\text{core}(\nu) \subseteq \text{span} \{P_1, \dots, P_N\}$,
- (ii) ν is an exact and generalized linear production game,
- (iii) there is a w^* -compact set $\Gamma_\alpha \subseteq \text{span} \{P_1, \dots, P_N\}$ such that $\nu(E) = \min_{m \in \Gamma_\alpha} m(E)$ for all E ,
- (iv) there is a function $g : \mathbb{R}^N \rightarrow \mathbb{R}$ concave and homogeneous of degree one such that $\nu(E) = g(P(E))$ for all E .

Then, (i) \Leftrightarrow (ii) \Leftrightarrow (iii) \Rightarrow (iv), with

$$g(x) = \min_{\{\chi : \chi \in R(\Gamma_\alpha)\}} \chi \cdot x$$

for all $x \in \mathbb{R}^N$, and

$$\text{core}(\nu) = \tilde{\Gamma}_\alpha = \text{co}(\Gamma_\alpha) = \{\chi \cdot P : \chi \in \text{co}(R(\Gamma_\alpha))\}. \quad (14)$$

Finally, if (iv) holds, there is an exact game $\nu_e = g_e(P)$, where

$$g_e(x) = \min_{\{\chi \in \partial g(0) : \chi \cdot P(\Omega) = g(P(\Omega))\}} \chi \cdot x$$

for each $x \in \mathbb{R}^N$, such that $\nu_e \geq \nu$ and $\text{core}(\nu_e) = \text{core}(\nu) \subseteq \text{span}\{P_1, \dots, P_N\}$.

Remark. Up to some obvious changes, Theorem 31, as well as Lemmas 29 and 30, still hold if P were a signed vector charge. In fact, we can replace (P_1, \dots, P_N) with the “enlarged” positive vector charge $(P_1^+, P_1^-, \dots, P_N^+, P_N^-)$ and then use the relation $\text{span}(P_1, \dots, P_N) \subseteq \text{span}(P_1^+, P_1^-, \dots, P_N^+, P_N^-)$. In this way all cores that are finite dimensional subsets of $sc(\Omega)$ are indeed covered by Theorem 31.

We close with a simple corollary of Theorem 31, which shows that exact games without “interior” linear sets cannot have finite-dimensional cores in $sc(\Omega)$.

Corollary 32 *Let ν be an exact game such that for all its linear sets A we have either $\nu(A) = 0$ or $\nu(A) = \nu(\Omega)$. If $\nu(\Omega) \neq 0$, then $\text{core}(\nu)$, when nonempty, is not a finite-dimensional subset of $sc(\Omega)$.*

Proof. Suppose $\text{core}(\nu) \neq \emptyset$ and suppose, *per contra*, that $\text{core}(\nu)$ is a finite-dimensional subset of $sc(\Omega)$. By Theorem 31, there is a function $g : \mathbb{R}^N \rightarrow \mathbb{R}$ concave and homogeneous of degree one such that $\nu(E) = g(P(E))$ for all E . Hence, given any diagonal set E_t , with $P(E_t) = tP(\Omega)$, it holds $\nu(E_t) = t\nu(\Omega)$. Therefore, E_t is linear and $\nu(E_t) \notin \{0, \nu(\Omega)\}$, a contradiction. ■

8 Concluding Remarks

1. Theorem 21 generalizes in several ways a well-known result of Billera and Raanan (1981), which establishes that cores of some measure games consist of linear combinations of measures (Corollary 2.6 p. 422).

First, their result requires the existence of a linear set A such that both $P(A) = 2^{-1}P(\Omega)$ and $\nu(A) = 2^{-1}\nu(\Omega)$. We only require A to be a linear and radial set.

Second, they require that $\nu \in pNA'$, the supnorm closure of polynomial functions of several non-atomic measures defined on a space isomorphic to $[0, 1]$ with its Borel sets (Aumann and Shapley (1974) p. 152). This topological structure is crucial for their results, and $g(P) \in pNA'$ if and only if g is continuous on $R(P)$. In contrast, we do not make any topological assumption, and our result holds for any measure game ν .⁵

Third, their Corollary 2.7 establishes the positivity of the coefficients of the linear combinations for nonatomic linear production games, a special class of measure games whose functions g are monotone. Our Theorem 21, instead, holds for any measure game having a monotone function g .

Finally, Theorem 21 follows from a subcalculus approach to the core and from a general linearity result for vector measures that put this result in a broader perspective. In particular, Proposition 25 and Corollary 26 are a dividend of this more general approach.

2. Corollary 23 extends some recent interesting results of Einy, Moreno, and Shitovitz (1999). Using different techniques, they prove (Theorem C) a special case of Corollary 23 for measure games whose function $g : R(P) \rightarrow \mathbb{R}$ is concave and continuous at $P(\Omega)$, rather than for general radially concave measure games, as we can do on the basis of our generalization of Billera and Raanan (1981).

3. Linear cores are very tractable objects. In fact, it is easy to check that to compute the linear core of a measure game $g(P)$ is enough to solve the following optimization problem in \mathbb{R}^N :

$$\begin{aligned} & \min_{(\alpha_1, \dots, \alpha_N) \in \mathbb{R}^N} \sum_{i=1}^N \alpha_i P_i(\Omega) \\ \text{s.t. } & \sum_{i=1}^N \alpha_i x_i \geq g(x_1, \dots, x_N) \quad \text{for all } (x_1, \dots, x_N) \in R(P). \end{aligned}$$

⁵Theorem 21 can be stated in the form used by Billera and Raanan (1981) for their Corollary 2.6, where ν is not necessarily a measure game. We prefer to use directly measure games, the most interesting class of games to which this class of results applies.

This problem is linear and involves finitely many variables – the coefficients $(\alpha_1, \dots, \alpha_N)$ – that appear in infinitely many constraint – the inequalities $\sum_{i=1}^N \alpha_i x_i \geq g(x_1, \dots, x_N)$ with $(x_1, \dots, x_N) \in R(P)$. Problems of this type are called semi-infinite linear problems and there is a large literature dealing with their theoretical and computational features (see, e.g., Goberna and Lopez, 1998). Since they involve only finitely many variables, computationally they are in general much more tractable than standard infinite programs and it is often possible to study them via their finite linear subprograms.

A Appendix

A.1 Proposition 6

The proof is based on the following lemma, which generalizes to bounded convex games a well-known properties of positive convex games.

Lemma 33 *Let $\nu : \Sigma \rightarrow \mathbb{R}$ be a bounded and convex game. Given any chain $\{E_i\}_{i \in I}$, there is $m \in \text{core}(\nu)$ such that $m(E_i) = \nu(E_i)$ for all $i \in I$.*

Proof. Given any Σ -measurable simple function $f : \Omega \rightarrow \mathbb{R}$, the Choquet integral $\int f d\nu$ is still well defined. Now, let $f, g : \Omega \rightarrow \mathbb{R}$ be any two Σ -measurable simple functions. Let $\Sigma_{f,g}$ be the smallest algebra that makes f and g measurable. As $\Sigma_{f,g}$ is finite, there is a (possibly zero) measure m on $\Sigma_{f,g}$ such that $\nu(E) \geq m(E)$ for all $E \in \Sigma_{f,g}$. Hence, $\nu - m$ is a positive convex game on $\Sigma_{f,g}$, and so, by a classic result of [6], $\int (f + g) d(\nu - m) \geq \int f d(\nu - m) + \int g d(\nu - m)$. In turn, this obviously implies $\int (f + g) d\nu \geq \int f d\nu + \int g d\nu$. We conclude that the Choquet integral $\int f d\nu$ is a superadditive functional on the vector space $B_0(\Sigma)$ of Σ -measurable simple functions.

Let Σ^* be a subalgebra of Σ on which there is a charge $m^* : \Sigma^* \rightarrow \mathbb{R}$ such that $m^* \in \text{core}(\nu|_{\Sigma^*})$. Since $\int f d\nu$ is a superadditive functional on $B_0(\Sigma)$, by the Hahn-Banach Theorem there is an extension $m : \Sigma \rightarrow \mathbb{R}$ of m^* such that $m \in \text{core}(\nu)$.

Consider now the chain $\{E_i\}_{i \in I}$. Let Σ_J be the algebra generated by a finite subchain $\{E_i\}_{i \in J}$. Let $m_J : \Sigma_J \rightarrow \mathbb{R}$ be the, possibly zero, charge on Σ_J such that $\nu(E) \geq m_J(E)$ for all $E \in \Sigma_J$. By well-known results (see, e.g., [7]), there exists $m' \in \text{core}(\nu - m_J)$ such that $m'(E_i) = (\nu - m_J)(E_i)$ for all $i \in J$. Hence, there is $m^* \in \text{core}(\nu|_{\Sigma_J})$ such that $m^*(E_i) = \nu(E_i)$ for

all $i \in J$. In turn, this implies the existence of an extension $m : \Sigma \rightarrow \mathbb{R}$ of m^* such that $m \in \text{core}(\nu)$.

Let $\Lambda_J = \{m \in \text{core}(\nu) : m(E_j) = \nu(E_j) \text{ for all } j \in J\}$. Since $\text{core}(\nu)$ is w^* -compact, the set Λ_J is w^* -compact. Moreover, by what we just proved, $\Lambda_J \neq \emptyset$. The collection $\{\Lambda_J\}_{\{J: J \subseteq I \text{ and } |J| < \infty\}}$ has the finite intersection property, and so its overall intersection is nonempty. Let m be an element of such intersection. We have $m \in \text{core}(\nu)$ and $m(E_i) = \nu(E_i)$ for all $i \in I$, as desired. ■

Proof of Proposition 6. By Lemma 33, there exists $m \in \text{core}(\nu)$ such that $m(E_1) = \nu(E_1)$ and $m(E_2) = \nu(E_2)$. This immediately implies that $m \in \partial\nu(E_1) \cap \partial\nu(E_2)$, and so $\partial\nu(E_1) \cap \partial\nu(E_2) \neq \emptyset$. As to the converse, suppose that $\partial\nu(E_1) \cap \partial\nu(E_2) \neq \emptyset$ for every $E_1 \subseteq E_2$. Let E and E' be any two sets of Σ . Let $m \in \partial\nu(E \cap E') \cap \partial\nu(E \cup E')$. Then, $\nu(E) \leq \nu(E \cup E') - m(E \cup E') + m(E)$ and $\nu(E') \leq \nu(E \cap E') - m(E \cap E') + m(E')$. By adding up we get $\nu(E) + \nu(E') \leq \nu(E \cup E') + \nu(E \cap E')$, as desired. Hence, ν is convex if and only if $\partial\nu(E_1) \cap \partial\nu(E_2) \neq \emptyset$. ■

A.2 Theorem 7

Let $B_1^+(\Sigma) = \{f \in B(\Sigma) : 0 \leq f \leq 1\}$, which are the ideal sets in the terminology of [2]. Given a bounded convex game ν , consider the functional $\nu^* : B(\Sigma) \rightarrow \mathbb{R}$ defined by

$$\nu^*(f) = \begin{cases} \int_0^{+\infty} \nu(f \geq t) dt & f \in B_1^+(\Sigma) \\ -\infty & f \notin B_1^+(\Sigma) \end{cases}$$

The integral $\int_0^{+\infty} \nu(f \geq t) dt$ is the Choquet integral of f w.r.t. ν . It is easy to check that, by Lemma 33, $\int_0^{+\infty} \nu(f \geq t) dt$ is a well defined Riemann integral. Again by Lemma 33, it is easy to check that ν^* is a proper concave function on $B(\Sigma)$. Moreover, $\nu^*(1_E) = \nu(E)$ for all $E \in \Sigma$.

Given $m \in \text{ba}(\Omega)$, let $L_m : B(\Sigma) \rightarrow \mathbb{R}$ be defined by $L_m(f) = \int f dm$. Let $\partial\nu^*(f)$ be the standard superdifferential of $\nu^* : B(\Sigma) \rightarrow \mathbb{R}$ at f . We show that $\partial\nu^*(1_E) = \{L_m : m \in \partial\nu(E)\}$ for all $E \in \Sigma$. Clearly, $\partial\nu^*(1_E) \subseteq \{L_m : m \in \partial\nu(E)\}$ for all $E \in \Sigma$. As to the converse inclusion, let $m \in \partial\nu(E)$. By definition, $\nu(A) - m(A) \leq \nu(E) - m(E)$ for all $A \in \Sigma$. Hence, for all $f \in B_1^+(\Sigma)$,

$$\nu^*(f) - L_m(f) = \int_0^1 (\nu - m)(f \geq t) dt \leq [\nu(E) - m(E)] \int_0^1 dt,$$

and so $L_m \in \partial\nu^*(1_E)$. This proves the converse inclusion, so that $\partial\nu^*(1_E) = \{L_m : m \in \partial\nu(E)\}$ for all $E \in \Sigma$. With a slight abuse of notation, we write $\partial\nu^*(1_E) = \partial\nu(E)$.

There is $f \in B_1^+(\Sigma)$ in a neighborhood of which (w.r.t. the norm topology) both ν_1^* and ν_2^* are bounded (e.g., $f = \alpha 1_\Omega$ for some $\alpha \in (0, 1)$). Then, by Theorem 20 of [22], for all $E \in \Sigma$ we have:

$$\begin{aligned} \partial(\nu_1 + \nu_2)(E) &= \partial(\nu_1 + \nu_2)^*(E) = \partial(\nu_1^* + \nu_2^*)(E) \\ &= \partial\nu_1^*(E) + \partial\nu_2^*(E) = \partial\nu_1(E) + \partial\nu_2(E), \end{aligned}$$

as desired. ■

A.3 Proposition 9

Consider condition (i). Let $x_0 \in R(P)$. Since g is concave on U , by a standard result in Convex Analysis (see Theorem 23.4 of [21]) there is a vector $\chi \in \mathbb{R}^N$ such that $g(x) \leq g(x_0) + \chi \cdot (x - x_0)$ for all $x \in U$. Hence, $\chi \cdot P \in \partial\nu(E)$ if $P(E) = x_0$.

Consider now condition (ii). Let K and W be respectively the cone and subspace generated by the convex set $R(P)$. Since $0 \in R(P)$, $W = K - K$ (see Theorem 2.7 of [21]). Define the function $g' : K \rightarrow \mathbb{R}$ by $g'(\lambda x) = \lambda g(x)$ with $x \in R(P)$ and $\lambda > 0$. The function g' is well-defined and it is superadditive and homogeneous of degree one on K . Define $g'' : W \rightarrow \mathbb{R}$ by $g''(w) = \sup \{g'(x) + g'(y) : x, y \in K \text{ and } x - y = w\}$. The function g'' as well is superadditive and homogeneous of degree one on W . Given any $x_0 \in R(P)$, let W_0 be the subspace of W generated by x_0 , i.e., $W_0 = \{\alpha x_0 : \alpha \in \mathbb{R}\}$. Define the linear functional $L_0 : W_0 \rightarrow \mathbb{R}$ by $L_0(\alpha x_0) = \alpha g(x_0)$ for all $\alpha \in \mathbb{R}$. If $\alpha \geq 0$, clearly $L_0(\alpha x_0) = g''(\alpha x_0)$ for all $w \in W_0$. If $\alpha < 0$, we have:

$$L_0(\alpha x_0) = \alpha g(x_0) = (-\alpha)(-g(x_0)) \geq (-\alpha)g''(-x_0) = g''(\alpha x_0).$$

By the Hahn-Banach Theorem, there exists a linear functional $L : W \rightarrow \mathbb{R}$ that extends L_0 on W and such that $L(w) \geq g''(w)$ for all $w \in W$. Since W is a subspace of \mathbb{R}^N , there exists a linear functional $L^* : \mathbb{R}^N \rightarrow \mathbb{R}$ that extends L on \mathbb{R}^N . Let $\chi^* \in \mathbb{R}^N$ such that $L^*(x) = \chi^* \cdot x$ for all $x \in \mathbb{R}^N$. Then, $\chi^* \cdot w \geq g''(w)$ for all $w \in W$ and $\chi^* \cdot x_0 = g(x_0)$. Hence, given any $x \in R(P)$, we have $g(x) - g(x_0) \leq \chi^* \cdot x - \chi^* \cdot x_0$, which implies $\chi^* \in \partial g(x_0)$. We conclude that $\partial g(x_0) \neq \emptyset$, as desired. ■

A.4 Theorem 14

Lemma 34 *Let $g(P) : \Sigma \rightarrow \mathbb{R}$ be a measure game with P countably additive. If g is lower Lipschitzian at 0 and $P(\Omega)$, there exists $\gamma > 0$ such that $\|m\|(E) \leq \gamma P^*(E)$ for all $E \in \Sigma$ and all $m \in \text{core}(\nu)$. In particular, all charges in $\text{core}(\nu)$ are countably additive.*

Remark. To prove that $\|m\|(\cdot) \leq \gamma P^*(\cdot)$ it suffices that P is finitely additive. In turn, this implies that m is strongly continuous.

Proof. Let $m \in \text{core}(\nu)$. By Theorem 10, $m \in \partial\nu(\Omega)$ and $m \in \partial\nu(\emptyset)$. Since $m \in \partial\nu(\Omega)$,

$$g(P(E)) - g(P(\Omega)) \leq -m(E^c) \quad (15)$$

for all $E \in \Sigma$. Moreover, since g is lower Lipschitzian at $P(\Omega)$, there exists $\gamma > 0$ and $\varepsilon_1 > 0$ such that

$$[g(P(E)) - g(P(\Omega))]^- \leq \gamma_1 \|P(E) - P(\Omega)\| \quad (16)$$

for all $E \in \Sigma$ such that $\|P(E) - P(\Omega)\| \leq \varepsilon_1$. Hence, Eqs. (15) and (16) imply that $m(E^c) \leq \gamma_1 \|P(E^c)\|$ for all $E \in \Sigma$ such that $\|P(E^c)\| \leq \varepsilon_1$. Since this holds for all $E \in \Sigma$, this implies that $m^+(E) \leq \gamma_1 \|P(E)\|$ for all $E \in \Sigma$ such that $\|P(E)\| \leq \varepsilon_1$. On the other hand, since $m \in \partial\nu(\emptyset)$, we have $g(P(E)) \leq m(E)$ for all $E \in \Sigma$, and, being g lower Lipschitzian at 0, there exists $\gamma_2 > 0$ and $\varepsilon_2 > 0$ such that $[g(P(E))]^- \leq \gamma_2 \|P(E)\|$ for all $E \in \Sigma$ such that $\|P(E)\| \leq \varepsilon_2$. Hence, $m^-(E) \leq \gamma_2 \|P(E)\|$ for all $E \in \Sigma$ such that $\|P(E)\| \leq \varepsilon_2$. Setting $\bar{\gamma} = \gamma_1 \vee \gamma_2$ and $\bar{\varepsilon} = \varepsilon_1 \wedge \varepsilon_2$, all this implies that $\|m\|(E) \leq 2\bar{\gamma} \|P(E)\|$ for all $E \in \Sigma$ such that $\|P(E)\| \leq \bar{\varepsilon}$. Since P is positive, there also exists $\gamma > 0$ such that

$$\|m\|(E) \leq \bar{\gamma} \|P(E)\| \leq \gamma P^*(E). \quad (17)$$

By the strong continuity of the component measures P_i , for each $E \in \Sigma$ there exists a partition $\{E_k\}_{k=1}^K$ of E in Σ such that $P^*(E_k) \leq \bar{\varepsilon}$ for each $k = 1, \dots, K$. Hence, by (17),

$$\|m\|(E) = \sum_{k=1}^K \|m\|(E_k) \leq \gamma \sum_{k=1}^K P^*(E_k) = \gamma P^*(E),$$

as desired. ■

Lemma 35 *Let $\nu = g(P) : \Sigma \rightarrow \mathbb{R}$ be a measure game and suppose P is countably additive. If there exists a linear set A such that g is lower semicontinuous at $P(A)$ and $P(A^c)$, then all charges in $\text{core}(\nu)$ are countably additive. Moreover, if $\text{core}(\nu) \neq \emptyset$, then g is continuous at $P(A)$ and $P(A^c)$.*

Proof. If $\text{core}(\nu) = \emptyset$, then obviously all charges in $\text{core}(\nu)$ are countably additive. Assume $\text{core}(\nu) \neq \emptyset$. We first prove that g is continuous at $P(A)$ and $P(A^c)$. Let $P(E_n) \rightarrow P(A)$. Since $\text{core}(\nu) \neq \emptyset$, $\nu(E_n) + \nu(E_n^c) \leq \nu(\Omega)$ for all $E_n \in \Sigma$. Hence,

$$\begin{aligned} & \limsup_n g(P(E_n)) + \liminf_n g(P(E_n^c)) \\ & \leq \limsup_n [g(P(E_n)) + g(P(E_n^c))] \leq g(P(\Omega)), \end{aligned}$$

and so, being g lower semicontinuous at $P(A)$ and $P(A^c)$,

$$\begin{aligned} \limsup_n g(P(E_n)) & \leq g(P(\Omega)) - \liminf_n g(P(E_n^c)) \\ & \leq g(P(\Omega)) - g(P(A^c)) = g(P(A)) \leq \liminf_n g(P(E_n)). \end{aligned}$$

Hence, $\lim_n g(P(E_n)) = g(P(A))$ and g is continuous at $P(A)$. By a similar method one can easily see that g is continuous at $P(A^c)$ as well.

To complete the proof it suffices to follow an argument similar to [2] p. 173. For, let $E_n \uparrow \Omega$ and let $m \in \text{core}(\nu)$. Then $P(E_n \cap A) \uparrow P(A)$ and $P(E_n \cap A^c) \uparrow P(A^c)$, and we can write:

$$\begin{aligned} \liminf_n m(A \cap E_n) & \geq \liminf_n g(P(A \cap E_n)) = g(P(A)) = g(P(\Omega)) - g(P(A^c)) \\ & = g(P(\Omega)) - \liminf_n g(P(A^c \cap E_n)) \\ & \geq m(\Omega) - \liminf_n m(A^c \cap E_n) = \limsup_n m(A \cap E_n), \end{aligned}$$

and so $\lim_n m(A \cap E_n) = m(A)$. On the other hand, a similar argument shows that $\lim_n m(A^c \cap E_n) = m(A^c)$. Hence, $\lim_n m(E_n) = m(\Omega)$, which implies that m is countably additive. ■

Proof of Theorem 14. In view of Lemma 35, all charges in $\text{core}(\nu)$ are countably additive. Let us now prove that all $m \in \text{core}(\nu)$ are absolutely continuous w.r.t. P^* . For, let E be such that $P^*(E) = 0$. Then, $P_i(E) = 0$

and $P_i(E^c) = 1$ for each $i = 1, \dots, N$ and so $m(E) \geq g(P(E)) = 0$ and $m(E^c) \geq g(P(E^c)) = \nu(\Omega) = m(\Omega)$. Hence,

$$0 \leq m(E) = m(\Omega) - m(E^c) \leq 0,$$

which implies $m(E) = 0$. Since $m \ll P^*$, by a variation of the Lebesgue Decomposition Theorem, there exist measures $\{m_i\}_{i=1}^N$ such that $m_i \ll P_i$ for each $i = 1, \dots, N$, and $m(E) = \sum_{i=1}^N m_i(E)$ for all E . Moreover, the measures $\{m_i\}_{i=1}^N$ are mutually singular and $\|m\|(E) = \sum_{i=1}^N \|m_i\|(E)$ for all E (see, e.g., Proposition 8.5.1 of [3]). By the Radon-Nikodym Theorem, there exists a Σ -measurable vector function $f = (f_1, \dots, f_N) : \Omega \rightarrow \mathbb{R}^N$ such that, for all $E \in \Sigma$, $m(E) = \sum_{i=1}^N \int_E f_i dP_i$.

Suppose $f : \Omega \rightarrow \mathbb{R}^N$ is such that for all $E \in \Sigma$ Eq. (4) holds. Then,

$$\begin{aligned} \int_E \frac{dm}{dP^*} dP^* &= m(E) = \sum_{i=1}^N \int_E f_i dP_i = \sum_{i=1}^N \int_E f_i \frac{dP_i}{dP^*} dP^* \\ &= \int_E \left(\sum_{i=1}^N f_i \frac{dP_i}{dP^*} \right) dP^*, \end{aligned}$$

and so $\sum_{i=1}^N f_i \frac{dP_i}{dP^*} = \frac{dm}{dP^*}$ P^* -a.e. Conversely, assume that $f : \Omega \rightarrow \mathbb{R}^N$ solves the relation (5). Then, for all $E \in \Sigma$,

$$\begin{aligned} m(E) &= \int \frac{dm}{dP^*} dP^* = \int_E \left(\sum_{i=1}^N f_i \frac{dP_i}{dP^*} \right) dP^* = \sum_{i=1}^N \int_E \left(f_i \frac{dP_i}{dP^*} \right) dP^* \\ &= \sum_{i=1}^N \int_E f_i dP_i, \end{aligned}$$

as desired.

Finally, suppose that g is lower Lipschitzian at 0 and $P(\Omega)$. By Lemma 34, all charges in $\text{core}(\nu)$ are countably additive. Moreover, since $m \ll P^*$, by the Radon-Nikodym Theorem there exists a Σ -measurable function f such that, for all sets E , $m(E) = \int_E f dP^*$ and $\|m\|(E) = \int_E |f| dP^*$. Set $A = \{|f| \geq c\}$ with $c > 0$. By Lemma 34, $\|m\| \leq \gamma P^*$. Hence,

$$\gamma P^*(|f| \geq c) \geq \|m\|(|f| \geq c) = \int_A |f| dP^* \geq c P^*(|f| \geq c),$$

which implies that $\gamma \geq c$ whenever $P^*(|f| \geq c) > 0$. Thus, $\|f\|_\infty \leq \gamma$. ■

A.5 Proposition 16

By Lemma 3, $\chi \cdot P \in \partial\nu(A)$ if and only if $\chi \in \partial g(P(A))$. On the other hand, by Theorem 10, $\chi \cdot P \in \text{core}(\nu)$ if and only if $\chi \cdot P \in \partial\nu(A) \cap \partial\nu(A^c)$. Hence, $\chi \cdot P \in \text{core}(\nu)$ if and only if $\chi \in \partial g(P(A)) \cap \partial g(P(A^c))$. This proves (6).

We now prove (7). It is trivially true if $\mathcal{L}\text{core}(\nu)$ is empty. Hence, assume that $\mathcal{L}\text{core}(\nu) \neq \emptyset$. Set $\dim(R(P)) = n$ and $\dim(\text{span}(\{P(A) : A \in \mathcal{A}\})) = k$. Clearly, $k \leq n \leq N$. Let $\{A_i\}_{i=1}^k \subseteq \mathcal{A}$ be such that the vectors $\{P(A_i)\}_{i=1}^k$ are linearly independent. Given $m \in \mathcal{L}\text{core}(\nu)$, we have $m(\cdot) = R(m) \cdot P(\cdot)$, where R is the canonical isomorphism (see Lemma 29 of Section 7). Hence, $\{R(m) : m \in \mathcal{L}\text{core}(\nu)\} \subseteq \mathbb{R}^N$ belongs to the affine space M defined by the linear equations $\xi \cdot P(A_i) = \nu(A_i)$ for $i = 1, \dots, k$. Since R is an isomorphism, $\dim(\mathcal{L}\text{core}(\nu)) \leq \dim(M)$. The dimension of M is equal to the dimension of the space M_0 defined by the homogeneous linear equations $\xi \cdot P(A_i) = 0$ for $i = 1, \dots, k$. As $M_0 = \text{span}(\{P(A) : A \in \mathcal{A}\})^\perp$, we conclude that $\dim(\mathcal{L}\text{core}(\nu)) \leq \dim(M) = n - k$, as desired.

Finally, being $P(\Omega) \neq 0$, $\dim(\text{span}(\{P(A) : A \in \mathcal{A}\})) \geq 1$, which proves the last inequality of (7). ■

A.6 Proposition 19

Since $2^{-1}P(\Omega)$ is the center of symmetry of $R(P)$, it is easy to see that $2^{-1}P(\Omega) \in \text{ri}(R(P))$ and that

$$\text{ri}(R(P)) = \bigcup_{t \in (0,1]} \left\{ t \frac{P(\Omega)}{2} + (1-t)R(P) \right\}. \quad (18)$$

Let $\bar{t} \in (0, 1)$. Suppose first that $\bar{t} \geq 1/2$. Setting $t' = 2(1 - \bar{t})$, we have $t' \in (0, 1)$. Moreover, it is easy to check that

$$t' \frac{P(\Omega)}{2} + (1-t')P(E) = \bar{t}P(E) + (1-\bar{t})P(E^c),$$

and so $\bar{t}P(E) + (1-\bar{t})P(E^c) \in \text{ri}(R(P))$. Next suppose that $\bar{t} < 1/2$. Setting $t' = 2\bar{t}$, we have $t' \in (0, 1)$. Moreover, it is easy to check that

$$t' \frac{P(\Omega)}{2} + (1-t')P(E^c) = \bar{t}P(E) + (1-\bar{t})P(E^c),$$

and so $\bar{t}P(E) + (1-\bar{t})P(E^c) \in \text{ri}(R(P))$. This completes the proof of the result. ■

A.7 Theorem 20

Assume that $m \in ca(\Omega)$. We first show that $P(E) = 0$ implies $m(E) = 0$ for all $E \in \Sigma$. In fact, consider the sets $E \cap A$ and $E \cap A^c$. We have $P(E \cap A) = P(E \cap A^c) = 0$, and so $P(A - E \cap A) = P(A)$ and $P(A^c - E \cap A^c) = P(A^c)$. By (8), this implies $m(A - E \cap A) = m(A)$ and $m(A^c - E \cap A^c) = m(A^c)$, so that $m(E \cap A) = m(E \cap A^c) = 0$, and we conclude that $m(E) = m(E \cap A) + m(E \cap A^c) = 0$.

Next we show that m is non-atomic. Let $m(E) \neq 0$. By what has been just proved, $P(E) \neq 0$. In particular, set $J = \{1 \leq i \leq N : P_i(E) > 0\}$ and $P^* = \{P_j\}_{j \in J}$. By Lyapunov Theorem there exists a partition E^1, B^1 of E such that $P^*(E^1) = P^*(B^1) = 2^{-1}P^*(E)$. If both $m(E^1) \neq 0$ and $m(B^1) \neq 0$, we are done. Suppose, in contrast, that either $m(E^1) = 0$ or $m(B^1) = 0$. W.l.o.g., suppose that $m(E^1) = m(E)$. Again by Lyapunov Theorem, there exists a partition E^2 and B^2 of E^1 such that $P^*(E^2) = P^*(B^2) = \frac{1}{2}P^*(E^1)$. If both $m(E^2) \neq 0$ or $m(B^2) \neq 0$, we are done. Suppose, in contrast, that either $m(E^2) = 0$ or $m(B^2) = 0$. W.l.o.g., assume that $m(E^2) = m(E^1)$. Proceeding in this way, either we find a set $B \subseteq E$ such that both $m(B) \neq 0$ and $m(E - B) \neq 0$, or we can construct a chain $\{E^n\}_{n \geq 1}$ such that $P^*(E^n) = 2^{-n}P^*(E)$ and $m(E^n) = m(E)$ for all $n \geq 1$. Hence, being $\bigcap_{n \geq 1} E^n \in \Sigma$, and $\bigcap_{n \geq 1} E^n \subseteq E$, we have $P^*(\bigcap_{n \geq 1} E^n) = 0$ and $m(\bigcap_{n \geq 1} E^n) = m(E) \neq 0$, a contradiction since we have $P^*(\bigcap_{n \geq 1} E^n) = 0$ iff $P(\bigcap_{n \geq 1} E^n) = 0$. Hence, there exists some set $B \subseteq E$ such that both $m(B) \neq 0$ and $m(E - B) \neq 0$, and so m is non-atomic.

Therefore, under both hypotheses on m , m is strongly continuous. Consequently, by the Lyapunov Theorem, the range $R(P, m)$ of $(P, m) : \Sigma \rightarrow \mathbb{R}^{N+1}$ is a convex subset of \mathbb{R}^{N+1} . Set $W = \text{span}(R(P, m))$ and let

$$R_{P(A)} = \{x \in \mathbb{R} : (P(A), x) \in R\},$$

where A is the set of Eq. (8). By Eq. (8), $R_{P(A)} = \{m(A)\}$. Hence, by Theorem 6.8 of [21], $(P(A), m(A)) \in \text{ri}(R)$. In turn, this implies that $(0, 1) \notin \text{span}(R)$. For, suppose to the contrary that $(0, 1) \in W$. Since $(P(A), m(A)) \in \text{ri}(R(P, m))$, there is $t > 0$ small enough so that $(P(A), m(A)) + t(0, 1) \in R(P, m)$. Since this contradicts Eq. (8), we conclude that $(0, 1) \notin W$.

By a standard separation theorem (see, e.g., Corollary 11.4.2 of [21]), there is $\pi \in \mathbb{R}^{N+1}$ and $\alpha \in \mathbb{R}$ such that, for all $y \in W$,

$$\pi \cdot y < \alpha < \pi \cdot (0, 1).$$

As $0 \in W$, $\alpha > 0$. Hence, $\pi \cdot (0, 1) > \alpha > 0$ implies $\pi_{N+1} > 0$. Moreover, since W is a vector space, for each $y \in W$ we have $\pi \cdot (\lambda y) < \alpha$ for all $\lambda > 0$. Then $\pi \cdot y \leq 0$, which implies $\pi \cdot y = 0$. Therefore, for all $(P(E), m(E)) \in R(P, m)$ we have $\pi_{N+1}m(E) + \sum_{i=1}^N \pi_i P_i(E) = 0$, and we conclude that $m \in \text{span} \{P_1, \dots, P_N\}$, with coefficients $\{-(\pi_i/\pi_{N+1})\}_{i=1}^N$.

It is easy to check that $R(P)$ is full dimensional if and only if the charges $\{P_i\}_{i=1}^N$ are linearly independent, which is the only case when the coefficients $\{\pi_i/\pi_{N+1}\}_{i=1}^N$ are unique.

Finally, if Eq. (10) holds, then $\{(x, -1) : x \in \mathbb{R}_+^N\} \cap W = \emptyset$. By now, it is easy to see that, by applying a standard separation result on these two closed and disjoint convex sets, we can find a vector $\pi \in \mathbb{R}^{N+1}$ with $\pi_i/\pi_{N+1} \leq 0$ for all $i = 1, \dots, N$, and such that $\pi_{N+1}m(E) + \sum_{i=1}^N \pi_i P_i(E) = 0$. Hence, $m \in \text{cone} \{P_1, \dots, P_N\}$. ■

A.8 Corollary 23

Proof. In view of Theorem 21, it suffices to prove that radially concave measure games admit radial and linear sets. Suppose first that $P(E) \neq P(E^c)$. By the Lyapunov Theorem, for each $\alpha \in (0, 1)$ there exists $E_\alpha \in \Sigma$ such that $P(E_\alpha) = \alpha P(E) + (1 - \alpha) P(E^c)$. Therefore, using (13),

$$\begin{aligned} \nu(E_\alpha) &= g(P(E_\alpha)) = g(\alpha P(E) + (1 - \alpha) P(E^c)) \\ &\geq \alpha g(P(E)) + (1 - \alpha) g(P(E^c)) \\ \nu(E_\alpha^c) &= g(P(E_\alpha^c)) = g((1 - \alpha) P(E) + \alpha P(E^c)) \\ &\geq (1 - \alpha) g(P(E)) + \alpha g(P(E^c)). \end{aligned}$$

Hence, $\nu(E_\alpha) + \nu(E_\alpha^c) \geq \nu(E) + \nu(E^c) = \nu(\Omega) \geq \nu(E_\alpha) + \nu(E_\alpha^c)$ because $\text{core}(\nu) \neq \emptyset$, and so each E_α is linear and radial. If $P(E) = P(E^c)$, then $P(E) = (1/2)P(\Omega)$ and so it is the center of symmetry of $R(P)$ and it belongs to $\text{ri}(R(P))$. Hence, E itself is linear and radial.

Finally, it is easy to check that $P(\Omega)/2$ is linear. In fact, $P(E_{1/2}) = 2^{-1}P(\Omega)$. ■

A.9 Proposition 24

Since g is concave on G , it is Lipschitzian relative to the compact set $R(P)$ (see, e.g., [21] Theorem 10.4), and so condition (ii) of Theorem 21 and Corollary 23 holds. Let $\delta(x)$ be the appropriate indicator function of $R(P)$ for

our setting, defined by

$$\delta(x) \equiv \delta(x \mid R(P)) = \begin{cases} 0 & x \in R(P) \\ -\infty & x \notin R(P) \end{cases}$$

Set $\tilde{g}(x) = g(x) + \delta(x)$ for each $x \in G$. Clearly, $\partial\tilde{g}(x) = \partial g_{|R(P)}(x)$ for all $x \in R(P)$. Since g is concave, by a well-known result (see, e.g., [21] Theorem 23.8), $\partial\tilde{g}(x_0) = \partial g(x_0) + \partial\delta(x_0)$ for all $x_0 \in G$. As well-known, $\partial\delta(x_0) = \{\chi \in \mathbb{R}^N : \chi \cdot x_0 \leq \chi \cdot x \text{ for all } x \in R(P)\}$. Let $x_0 \in ri(R(P))$ and let $w \in W$, where $W = span(R(P))$. There is $\varepsilon > 0$ such that $x_0 + \varepsilon w \in R(P)$, and so $\chi \cdot w \geq 0$ for all $\chi \in \partial\delta(x_0)$. Since W is a vector subspace, this implies $\chi \cdot w = 0$ for all $\chi \in \partial\delta(x_0)$, which in turn implies that $\partial\delta(x_0) \subseteq W^\perp$. Since the converse inclusion is obvious, we conclude that $\partial\delta(x_0) = W^\perp$.

Putting everything together, we have $\partial g_{|R(P)}(x_0) = \partial g(x_0) + W^\perp$ for all $x_0 \in ri(R(P))$. Hence, given any $\chi \in \partial g_{|R(P)}(x_0)$, there is $\chi' \in \partial g(x_0)$ such that $\chi \cdot P = \chi' \cdot P$. Since $\partial g(x_0) \subseteq \partial g_{|R(P)}(x_0)$, this implies that $\{\chi \cdot P : \chi \in \partial g_{|R(P)}(P(A))\} = \{\chi \cdot P : \chi \in \partial g(P(A))\}$. A simple application of Corollary 23 now completes the proof. ■

A.10 Proposition 25

Suppose $core(\nu) \neq \emptyset$. This implies that $\partial g(P(A)) \cap \partial g(P(A^c)) \neq \emptyset$ because, by Theorem 21, $core(\nu) = \{\chi \cdot P : \chi \in \partial g(P(A)) \cap \partial g(P(A^c))\}$. Moreover, since $P(A) \in ri(R(P))$, by a well-known result of Convex Analysis, $[\nabla g(P(A)) - \chi] \cdot w = 0$ for each $\chi \in \partial g(P(A))$ and each $w \in W$, where $W = span(R(P))$. Hence, $\nabla g(P(A)) \cdot P(\cdot) = \chi \cdot P(\cdot)$ for each $\chi \in \partial g(P(A))$, and so, by Lemma 3, $\nabla g(P(A)) \cdot P(\cdot) \in \partial\nu(A)$. Since in [12] it is proved that $\delta\nu(\cdot; A) = \nabla g(P(A)) \cdot P(\cdot)$, we then have $\delta\nu(\cdot; A) \in \partial\nu(A)$. By Theorem 12, $core(\nu) = \{\delta\nu(\cdot; A)\}$, as desired. Next, suppose that g is differentiable and superdifferentiable at both $P(A)$ and $P(A^c)$. By proceeding as before, it can be shown that $\delta\nu(\cdot; A^c) \in \partial\nu(A^c)$. Hence, by Theorem 12, $core(\nu) \subseteq \{\delta\nu(\cdot; A^c)\}$, and so $core(\nu) \neq \emptyset$ implies $\delta\nu(\cdot; A) = \delta\nu(\cdot; A^c)$, i.e., $\nabla g(P(A)) = \nabla g(P(A^c))$. As to the converse, since $\delta\nu(\cdot; A) \in \partial\nu(A)$ and $\delta\nu(\cdot; A^c) \in \partial\nu(A^c)$, the equality $\delta\nu(\cdot; A) = \delta\nu(\cdot; A^c)$ implies $\partial\nu(A) \cap \partial\nu(A^c) \neq \emptyset$. Then, by Theorem 10, $core(\nu) \neq \emptyset$. ■

A.11 Lemmas 28, 29, and 30

Proof of Lemma 28. (i) is obvious. As to (ii), let $m_1, m_2 \in \tilde{\Gamma}_\alpha$ and let $E \in \Sigma$. Then, there exist $m'_1, m'_2 \in \Gamma_\alpha$ such that $m_1(E) \geq m'_1(E)$ and $m_2(E) \geq m'_2(E)$. Hence, for each $t \in (0, 1)$,

$$tm_1(E) + (1-t)m_2(E) \geq tm'_1(E) + (1-t)m'_2(E) \geq \min\{m'_1(E), m'_2(E)\}.$$

The set $\tilde{\Gamma}_\alpha$ is therefore convex. It is also immediate to check that $\tilde{\Gamma}_\alpha$ is closed. To show that it is w^* -compact, by the Alaoglu Theorem it is enough to prove that $\tilde{\Gamma}_\alpha$ is norm bounded. Given $E \in \Sigma$, for each $m \in \tilde{\Gamma}_\alpha$ there are $m', m'' \in \Gamma_\alpha$ such that $m'(E) \leq m(E) \leq m''(E)$ (consider E and E^c). Hence, $\sup_{m \in \tilde{\Gamma}_\alpha} |m(E)| = \sup_{m \in \Gamma_\alpha} |m(E)| < \infty$ and $\tilde{\Gamma}_\alpha$ is setwise bounded. By a variation of the Nikodym Boundedness Theorem (see, e.g., [8] p. 80), $\tilde{\Gamma}_\alpha$ is then norm bounded. To prove (iii), observe that $\nu(E) = \min_{m \in \tilde{\Gamma}_\alpha} m(E)$ is well defined because $\tilde{\Gamma}_\alpha$ is w^* -compact. Clearly, $\tilde{\Gamma}_\alpha \subseteq \text{core}(\nu)$. As to the converse, let $m \in \text{core}(\nu)$. By construction, for each E there is $m' \in \tilde{\Gamma}_\alpha$ such that $m'(E) \leq m(E)$. On the other hand, by the definition of m -closure, there is $m'' \in \Gamma_\alpha$ such that $m'(E) \geq m''(E)$, so that $m(E) \geq m''(E)$. Hence, $m \in \tilde{\Gamma}_\alpha$, and we conclude that $\tilde{\Gamma}_\alpha = \text{core}(\nu)$. The uniqueness of ν is obvious. ■

Proof of Lemma 29. We first show that R is well defined. Let $\chi, \chi' \in \mathbb{R}^N$ be such that $\chi \cdot P = \chi' \cdot P$. Then, $(\chi - \chi') \cdot P(E) = 0$ for all $E \in \Sigma$, and so $(\chi - \chi') \cdot P \in \text{span}(R(P))^\perp$. Hence, $\pi(\chi) - \pi(\chi') = \pi(\chi - \chi') = 0$, which implies $\pi(\chi) = \pi(\chi')$. It is easy to check that R is a linear isomorphism. Moreover, $\text{span}\{P_1, \dots, P_N\}$ with the relative w^* -topology is a finite-dimensional topological vector space. Hence, by a standard result (see, e.g., [24] p. 79), R is w^* -continuous. ■

Proof of Lemma 30. By using the canonical isomorphism R , we can write $m(\cdot) = R(m) \cdot P(\cdot)$ for each $m \in \Gamma_\alpha$. Consider the function $g(x) = \min_{m \in \Gamma_\alpha} R(m) \cdot x$ for all $x \in \mathbb{R}^N$ and the associated measure game $\nu = g(P)$, which is

$$\nu(\cdot) = \min_{m \in \Gamma_\alpha} R(m) \cdot P(\cdot).$$

Clearly, ν is exact and, by Lemma 28, $\text{core}(\nu) = \tilde{\Gamma}_\alpha$.

On the other hand, the canonical map R is continuous on the w^* -compact set Γ_α , which is metrizable since it is a finite-dimensional subset of $ba(\Omega)$.

Hence, by proceeding as in the example of generalized linear production game, we have

$$\begin{aligned} \text{core}(\nu) &= \{\chi \cdot P : \chi \in \text{co}(R(m) : R(m) \cdot P(\Omega) = \nu(\Omega)) = \alpha\} \\ &= \{\chi \cdot P : \chi \in \text{co}(R(m) : m \in \Gamma_\alpha)\} = \text{co}(R(m) \cdot P : m \in \Gamma_\alpha) \\ &= \text{co}(\Gamma_\alpha). \end{aligned}$$

We conclude that $\tilde{\Gamma}_\alpha = \text{co}(\Gamma_\alpha)$, as desired. ■

A.12 Theorem 31

If we set $\Gamma_\alpha = \text{core}(\nu)$, clearly (i) implies (iii). As to (iii) \Rightarrow (i), we have $\nu(E) = \min_{m \in \Gamma_\alpha} m(E)$ for all $E \in \Sigma$, and so $\nu(E) = \min_{m \in \tilde{\Gamma}_\alpha} m(E)$ for all E . By point (iii) of Lemma 28, $\text{core}(\nu) = \tilde{\Gamma}_\alpha$. Hence, ν is exact and, by Lemma 30, $\text{core}(\nu) \subseteq \text{span}\{P_1, \dots, P_N\}$. This proves that (iii) implies (i).

Set $g(x) = \min_{\chi \in R(\Gamma_\alpha)} \chi \cdot x$ for all $x \in \mathbb{R}^N$. The function $g : \mathbb{R}^N \rightarrow \mathbb{R}$ is well defined because $R(\Gamma_\alpha)$ is a compact subset of \mathbb{R}^N . Clearly, it is also concave and homogeneous of degree one, and, by point (iii), $\nu = g(P)$ since $\Gamma_\alpha = \{\chi \cdot P : \chi \in R(\Gamma_\alpha)\}$. Therefore, (iii) implies (iv), and (ii) and (iii) are equivalent.

As Lemma 30 immediately implies Eq. (14), to complete the proof it only remains to prove the last part of the theorem. Hence, assume that (iv) holds and set $C_e = \{\chi \in \partial g(0) : \chi \cdot P(\Omega) = g(P(\Omega))\}$. Define $g_e(x) = \min_{\chi \in C_e} \chi \cdot x$ for all $x \in \mathbb{R}^N$ and $\nu_e(\cdot) = g_e(P(\cdot))$. By construction, g_e is concave and homogeneous of degree one, and $\nu_e \geq \nu$. Since all diagonal sets are linear for both ν and ν_e , by Corollary 23,

$$\begin{aligned} \{\chi \cdot P : \chi \in C_e\} &= \{\chi \cdot P : \chi \in \partial g(0) \cap \partial g(P(\Omega))\} \\ &= \text{core}(\nu) \supseteq \text{core}(\nu_e) \supseteq \{\chi \cdot P : \chi \in C_e\}, \end{aligned}$$

and so $\text{core}(\nu_e) = \text{core}(\nu)$. ■

References

- [1] R. Aumann and J. Dreze, Cooperative Games with Coalition Structures, *International Journal of Game Theory* 3, 217-237, 1974.

- [2] R. Aumann and L. Shapley, *Values of Non-Atomic Games*, Princeton University Press, Princeton, 1974.
- [3] K.P.S.M. Bhaskara Rao, M. Bhaskara Rao, *Theory of Charges*, Academic Press, New York, 1983.
- [4] L. J. Billera and J. Raanan, Cores of Nonatomic Linear Production Games, *Mathematics of Operations Research* 8, 420-423, 1981.
- [5] E. Bolker, A Class of Convex Bodies, *Transactions of the American Mathematical Society* 145, 323-345, 1969.
- [6] G. Choquet, Theory of Capacities, *Annales de l'Institut Fourier* 5, 131-295, 1953.
- [7] F. Delbaen, Convex Games and Extreme Points, *Journal of Mathematical Analysis and Applications* 45, 210-233, 1974.
- [8] J. Diestel, *Sequences and Series in Banach Spaces*, Springer, New York, 1984.
- [9] N. Dunford and J.T. Schwartz, *Linear Operators, Part I: General Theory*, Wiley-Interscience, London, 1954.
- [10] E. Einy, D. Moreno and B. Shitovitz, The Core of a Class of Non-Atomic Games which Arise in Economic Applications, *International Journal of Game Theory* 28, 1-14, 1999.
- [11] L.G. Epstein, A Definition of Uncertainty Aversion, *Review of Economic Studies* 66, 579-608, 1999.
- [12] L. G. Epstein and M. Marinacci, The Core of Large Differentiable TU Games, *Journal of Economic Theory*, forthcoming.
- [13] P. C. Fishburn, On Harsanyi's Utilitarian Cardinal Welfare Theorem, *Theory and Decision* 17, 21-28, 1984.
- [14] S. Fujishige, *Submodular Functions and Optimization*, North-Holland, Amsterdam, 1991.
- [15] M. A. Goberna and M. A. Lopez, *Linear Semi-Infinite Optimization*, Wiley, New York, 1998.

- [16] S. Hart and A. Neyman, Values of Non-Atomic Vector Measure Games, *Journal of Mathematical Economics* 17, 31-40, 1988.
- [17] J. B. Hiriart-Urruty and C. Lemarechal, *Convex Analysis and Minimization Algorithms I*, Springer-Verlag, New York, 1993.
- [18] M. Marinacci, A Uniqueness Theorem for Convex-Ranged Measures, *Decisions in Economics and Finance* 23, 121-132, 2000.
- [19] P. Mongin, Consistent Bayesian Aggregation, *Journal of Economic Theory* 66, 313-351, 1995.
- [20] G. Owen, On the Core of Linear Production Games, *Mathematical Programming* 9, 358-370, 1975.
- [21] R. T. Rockafellar, *Convex Analysis*, Princeton University Press, Princeton, New Jersey, 1970.
- [22] R. T. Rockafellar, *Conjugate Duality and Optimization*, SIAM, Philadelphia, 1974.
- [23] D. Schmeidler, Cores of Exact Games, *Journal of Mathematical Analysis and Applications* 40, 214-225, 1972.
- [24] F. Trèves, *Topological Vector Spaces, Distributions and Kernels*, Academic Press, New York, 1967.

INTERNATIONAL CENTRE FOR ECONOMIC RESEARCH
APPLIED MATHEMATICS WORKING PAPER SERIES

1. Luigi Montrucchio and Fabio Privileggi, "On Fragility of Bubbles in Equilibrium Asset Pricing Models of Lucas-Type," *Journal of Economic Theory*, forthcoming (Icer WP 2001/5).
2. Massimo Marinacci, "Probabilistic Sophistication and Multiple Priors," *Econometrica*, forthcoming (Icer WP 2001/8).
3. Massimo Marinacci and Luigi Montrucchio, "Subcalculus for Set Functions and Cores of TU Games," April 2001 (Icer WP 2001/9).